



## Charge Radii of B and D Mesons in a Quark Model with Two Loop Static Potential

K. K. Pathak<sup>1\*</sup>, Namita Sharma Bordoloi<sup>2</sup> and D. K. Choudhury<sup>3</sup>

<sup>1</sup>Department of Physics, Arya Vidyapeeth College, Guwahati-781016, India.

<sup>2</sup>Department of Physics, Cotton College, Guwahati-781001, India.

<sup>3</sup>Department of Physics, Gauhati University, Guwahati-781014, India.

### Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/PSIJ/2015/18709

#### Editor(s):

- (1) Felix A. Buot, Center of Computational Materials Science, George Mason University, University Drive, Fairfax, Virginia, USA.
- (2) Christian Brosseau, Distinguished Professor, Department of Physics, Universit de Bretagne Occidentale, France.

#### Reviewers:

- (1) Anonymous, Carnegie Mellon University, USA.
- (2) Anonymous, University of Missouri, USA.
- (3) Anonymous, Cheju National University, South Korea.
- (4) Anonymous, Benha University, Egypt.

Complete Peer review History: <http://sciencedomain.org/review-history/10192>

### Original Research Article

Received: 6<sup>th</sup> May 2015  
Accepted: 24<sup>th</sup> June 2015  
Published: 15<sup>th</sup> July 2015

## Abstract

We study the effects of two loop static potential on the charge radii of heavy light flavoured mesons in an Improved QCD potential model. The estimated value of charge radii of heavy-light mesons in V-scheme is then compared with other available data and found to be within the limit of other result.

*Keywords:* Two loop static potential; Dalgarno method and charge radii.

\*Corresponding author: E-mail: [kkingkar@gmail.com](mailto:kkingkar@gmail.com)

## 1 INTRODUCTION

The study of hadron wavefunctions and their phenomenology is one of the important topics in QCD [1] and one of the reliable method of studying the physical properties of low energy QCD is the lattice formulation of gauge theory. But lattice QCD is limited to the Euclidean formulation of QCD and cannot be applied in Minkowski space time to simulate high energy reactions in which the particles are inherently moving near the light cone. It is also difficult to understand from simulations the important QCD mechanisms that lead to colour confinement. The only other way to proceed in the non perturbative regime of QCD is by inventing and using phenomenological models that capture the most important features of strong QCD. Since the classical work of De Rujula et al [2], such studies in non-relativistic potential approach have been made in several models [3, 4, 5, 6, 7, 8, 9, 10] with considerable success. Inspired by the success of this approach, the present authors have been pursuing a specific QCD Inspired Quark model [11, 12, 13, 14, 15] which has been developed in various stages : in earlier version [11, 12, 13, 14] significant confinement effects could not be introduced due to perturbative constraints, while in the later version [15] it could be incorporated.

The structure of the mesons can be learned by probing them with virtual photons in scattering

experiment. The form factors of mesons not only provide a testing ground for QCD-inspired models, but also are important in many areas of particle and nuclear physics, including charge radii, parity-violating experiments, and many others. Present paper reports the results of charge radii of heavy light mesons within the improved version of the model, where the linear part of the Cornell Potential  $V(r) = -\frac{4}{3}\frac{\alpha_s}{r} + br + c$ , is treated as perturbation and Coulombic part as parent to obtain an wavefunction by using Dalgarno method of perturbation[11, 13]. The corresponding results for the Isgur-Wise function and its slope and curvature have already been reported in [15].

In section 2 , we discuss the formalism , while in section 3 we summarise the results. Section 4 contains summary and conclusion .

## 2 FORMALISM

### 2.1 The Improved QCD Inspired Quark Model

The essential features of the improved model has already been reported in ref [15]. With the linear part " $br+c$ " of the Cornell potential as perturbation and coulombic part " $-\frac{4}{3}\frac{\alpha_s}{r}$ " as parent, the wave function in this model is given by

$$\psi_{rel+conf}(r) = \frac{N'}{\sqrt{\pi a_0^3}} \left( C'(c) - \frac{1}{2} \mu b a_0 r^2 \right) \left( \frac{r}{a_0} \right)^{-\epsilon} e^{-\frac{r}{a_0}} \quad (2.1)$$

where

$$N' = \frac{2^{\frac{1}{2}}}{\left[ 2^{2\epsilon} (\Gamma(3-2\epsilon) \{C'(c)\}^2 - \frac{1}{4} \mu b a_0^3 \Gamma(5-2\epsilon) C'(c) + \frac{1}{64} \mu^2 b^2 a_0^6 \Gamma(7-2\epsilon)) \right]^{\frac{1}{2}}} \quad (2.2)$$

is the normalisation constant;  $\epsilon$  and  $a_0$  are defined by

$$\epsilon = 1 - \sqrt{1 - \frac{4}{3} \alpha_s} \quad (2.3)$$

and

$$a_0 = \left( \frac{4}{3 \mu \alpha_{MS}} \right)^{-1} \quad (2.4)$$

respectively, with

$$C'(c) = 1 + c A_0 \sqrt{\pi a_0^3} \quad (2.5)$$

$A_0$  being the undetermined factor appearing in the series solution of the Schrödinger equation and  $\mu$  being the reduced mass.

From equation (5), if  $A_0$  and/or  $c$  is set equal to zero, the effect of  $c$  disappears in the formalism as was the case for [11, 12, 13, 14, 16]. From equations (1) and (2), it is also evident that in the limit  $b \rightarrow 0$ , the effect of  $C'(c)$  and hence of  $c$  completely disappears in the wave function.

## 2.2 Two Loop Static Potential in V Scheme

Traditionally, the strength of the quark gluon interaction is characterised by the coupling constant  $\alpha_{\overline{MS}}(q^2)$  defined by convention in a particular dimensional regularisation scheme such as the  $\overline{MS}$  regularisation scheme. V-scheme is a physically based alternative to this usual scheme. Indeed in lattice calculation,  $\alpha_V$

is regarded as the better expansion parameter. From the points of convergence and scale ambiguities as well, V-scheme is advocated in recent literature. The relationship of  $\alpha_V(\frac{1}{r^2})$  to the conventional coupling constant in  $\overline{MS}$  scheme at the two loop level has been reported by Peter[17, 18]. Later, Schröder has improved the work where the error in the gluonic sector has been corrected [19, 20].

Thus, the V-scheme ( V denoting potential ) [17, 18, 19, 20] defines the QCD coupling constant in terms of a potential, taking into account the higher order effects of QCD which are then expressed as a power series in the coupling constant  $\alpha_{\overline{MS}}$  in  $\overline{MS}$  scheme. In the V scheme  $\alpha_V$  represents the effective coupling constant which incorporates the entire radiative corrections into its definition. For completeness, we write the relationship between  $\alpha_V$  and  $\alpha_{\overline{MS}}$  at the NNLO level which is given by

$$\alpha_V \left( \frac{1}{r^2} \right) = \alpha_{\overline{MS}} \left( 1 + \frac{\alpha_{\overline{MS}}}{4\pi} (a_1 + \beta_0 L) + \frac{\alpha_{\overline{MS}}^2}{16\pi^2} \left[ a_2 + \beta_0^2 \left( L^2 + \frac{\pi^2}{3} \right) + (\beta_1 + 2\beta_0 a_1) L \right] + O(\alpha_{\overline{MS}}^3) \right) \quad (2.6)$$

where

$$L = 2 \ln(\tilde{\mu} r \exp \gamma), \quad (2.7)$$

$$\beta_0 = \frac{11}{3} C_A - \frac{4}{3} T_F n_f \quad (2.8)$$

and

$$\beta_1 = \frac{34}{3} C_A^2 - 4 C_F T_F n_f - \frac{20}{3} C_A T_F n_f \quad (2.9)$$

are the first two terms of the renormalisation group beta function,  $\tilde{\mu}$  being the scale parameter. The one-loop and two-loop constants with gauge independent coefficients are given by

$$a_1 = \frac{31}{9} C_A - \frac{20}{9} T_F n_f, \quad (2.10)$$

and

$$a_2 = \left( \frac{4343}{162} + 4\pi^2 - \frac{\pi^4}{4} + \frac{22}{3} \rho(3) \right) C_A^2 - \left( \frac{55}{3} - 16\rho(3) \right) C_F T_F n_f + \frac{400}{81} T_F^2 n_f^2 - \left( \frac{1798}{81} + \frac{56}{3} \rho(3) \right) C_A T_F n_f \quad (2.11)$$

### 2.3 Form Factor and Charge Radii.

The elastic charge form factor for a charged system of point quarks has the form [21, 22]

$$e \{F(Q^2)\} = \sum_{i=1}^2 \frac{e_i}{Q_i} \int_0^\infty r |\Psi(r)|^2 \sin(Q_i r) dr \quad (2.12)$$

where  $e_i$  is the charge of the  $i^{th}$  quark/antiquark and

$$Q_i = \frac{\sum_{j \neq i} m_j Q}{\sum_{i=1}^2 m_i} \quad (2.13)$$

$Q_i$  describes how the virtuality  $Q$  is shared between the quark antiquark pair of the meson. For a pair of same flavour (light-light or heavy-heavy), it is shared equally ( $Q_i = \frac{Q}{2}$ ), while for heavy-light, the entire virtuality is carried by the light quark. Using equation (1) in (12) and integrating over  $r$

$$e \{F(Q^2)\} = \frac{N'^2}{a_0 2^{-2\epsilon}} \sum_{i=1}^2 \frac{e_i}{Q_i} \left[ C'^2 \Gamma(2-2\epsilon) \sin[(2-2\epsilon)\theta_i] \left(1 + \left(\frac{1}{4} a_0^2 Q_i^2\right)\right)^{(\epsilon-1)} - \frac{1}{4} \mu b a_0^3 C' \Gamma(4-2\epsilon) \sin[(4-2\epsilon)\theta_i] \left(1 + \left(\frac{1}{4} a_0^2 Q_i^2\right)\right)^{(\epsilon-2)} + \frac{1}{64} \mu^2 b^2 a_0^6 \Gamma(6-2\epsilon) \sin[(6-2\epsilon)\theta_i] \left(1 + \left(\frac{1}{4} a_0^2 Q_i^2\right)\right)^{(\epsilon-3)} \right] \quad (2.14)$$

where

$$\theta_i = \sin^{-1} \frac{Q_i}{\left(\frac{4}{a_0^2} + Q_i^2\right)^{\frac{1}{2}}} \quad (2.15)$$

In evaluating (14) we use the approximation in (15) to be

$$\sin^{-1} x \approx x + \frac{x^3}{6} + \frac{3x^5}{40} \quad (2.16)$$

with

$$x = \frac{Q_i}{\left(\frac{4}{a_0^2} + Q_i^2\right)^{\frac{1}{2}}} \quad (2.17)$$

We also use

$$\sin y = y - \frac{y^3}{3!} + \frac{y^5}{5!} \quad (2.18)$$

with

$$y = (2-2\epsilon)\theta_i \quad (2.19)$$

As a result of these improved approximations, the expression for  $e \{F(Q^2)\}$  in equation (14) becomes

$$e \{F(Q^2)\} = \frac{N_1^2}{a_0 2^{-2\epsilon}} \sum_{i=1}^2 e_i \left[ \left(1 + \frac{1}{4} a_0^2 Q_i^2\right)^{(\epsilon-1)} \left( (2-2\epsilon) X_i - \frac{1}{6} (2-2\epsilon)^3 Q_i^2 X_i^3 + \frac{1}{120} (2-2\epsilon)^5 Q_i^4 X_i^5 \right) C'^2 \Gamma(2-2\epsilon) - \frac{1}{4} \mu b a_0^3 \left(1 + \frac{1}{4} a_0^2 Q_i^2\right)^{(\epsilon-2)} \left( (4-2\epsilon) X_i - \frac{1}{6} (4-2\epsilon)^3 Q_i^2 X_i^3 + \frac{1}{120} (4-2\epsilon)^5 Q_i^4 X_i^5 \right) C' \Gamma(4-2\epsilon) + \frac{1}{64} \mu^2 b^2 a_0^6 \left(1 + \frac{1}{4} a_0^2 Q_i^2\right)^{(\epsilon-3)} \left( (6-2\epsilon) X_i - \frac{1}{6} (6-2\epsilon)^3 Q_i^2 X_i^3 + \frac{1}{120} (6-2\epsilon)^5 Q_i^4 X_i^5 \right) \Gamma(6-2\epsilon) \right] \quad (2.20)$$

where

$$X_i = \frac{3Q_i^4}{40} \left( \frac{4}{a_0^2} + Q_i^2 \right)^{-\frac{5}{2}} + \frac{Q_i^2}{6} \left( \frac{4}{a_0^2} + Q_i^2 \right)^{-\frac{3}{2}} + \left( \frac{4}{a_0^2} + Q_i^2 \right)^{-\frac{1}{2}}, \quad (2.21)$$

$N_1$  being the normalization constant,

$$N_1 = 2^{\frac{1}{2}} \left( 2^{2\epsilon} \left( (2 - 2\epsilon) C'^2 \Gamma(2 - 2\epsilon) - \frac{1}{4} \mu b a_0^3 (4 - 2\epsilon) C' \Gamma(4 - 2\epsilon) + \frac{1}{64} \mu^2 b^2 a_0^6 (6 - 2\epsilon) \Gamma(6 - 2\epsilon) \right) \right)^{-\frac{1}{2}}. \quad (2.22)$$

The average charge radii square for the mesons are obtained from the relation

$$\langle r^2 \rangle = -6 \frac{dF(Q^2)}{dQ^2} \Big|_{Q^2=0}. \quad (2.23)$$

Using (20) in (23), one obtains for the mesons having quark masses  $m_i$  and  $m_j$ , and charges  $e_i$  and  $e_j$  respectively ,

$$\begin{aligned} \langle r^2 \rangle = a_0^2 \left[ e_i \left( 1 + \frac{m_i}{m_j} \right)^{-2} + e_j \left( 1 + \frac{m_j}{m_i} \right)^{-2} \right] & \left( C'^2 \Gamma(2 - 2\epsilon) \left( \frac{(2 - \epsilon)}{2} + \frac{(2 - \epsilon)^3}{4} + \right. \right. \\ & \left. \left. \frac{3(2 - \epsilon)(\epsilon - 1)}{2} \right) - \frac{1}{4} \mu b a_0^3 C' \Gamma(4 - 2\epsilon) \left( \frac{(4 - 2\epsilon)}{2} + \frac{(4 - 2\epsilon)^3}{4} + \frac{3(4 - 2\epsilon)(\epsilon - 2)}{8} \right) \right) \\ + \frac{1}{64} \mu^2 b^2 a_0^6 \Gamma(6 - 2\epsilon) & \left( \frac{(6 - 2\epsilon)}{2} + \frac{(6 - 2\epsilon)^3}{4} + \frac{3(6 - 2\epsilon)(\epsilon - 3)}{128} \right) \left( (2 - 2\epsilon) C'^2 \Gamma(2 - 2\epsilon) - \right. \\ & \left. \frac{1}{4} \mu b a_0^3 (4 - 2\epsilon) C' \Gamma(4 - 2\epsilon) + \frac{1}{64} \mu^2 b^2 a_0^6 (6 - 2\epsilon) \Gamma(6 - 2\epsilon) \right)^{-1} \end{aligned} \quad (2.24)$$

### 3 RESULTS AND DISCUSSION

#### 3.1 Relationship between $\alpha_{\overline{MS}}$ and $\alpha_V$

Taking number of quark flavour  $n_f = 4$  and  $n_f = 5$  with the corresponding value of  $\alpha_{\overline{MS}}$  at the scale of the c- and b- quark mass,  $\alpha_{\overline{MS}}(m_c) = 0.39$  and  $\alpha_{\overline{MS}}(m_b) = 0.22$  respectively [23], the value of  $\alpha_V(\frac{1}{r^2})$  for three choices of  $\tilde{\mu}$  are calculated[14] and shown in Table 1.

Table 1: Values of  $\alpha_V(\frac{1}{r^2})$  for different choices of  $\tilde{\mu}$  and the percentage increase

Choices of $\tilde{\mu}$	$\tilde{\mu} = \frac{1}{r}$	%	$\tilde{\mu} = \frac{1}{r} \exp\left(-\gamma_E - \frac{\alpha_1}{2\beta_0}\right)$	%	$\tilde{\mu} = \frac{1}{r^2} \exp(-\gamma_E)$	%
$\alpha_{\overline{MS}}(m_c) = 0.39,$ $n_f = 4$	0.693	77.7	0.625	66.9	0.604	54.9
$\alpha_{\overline{MS}}(m_b) = 0.22,$ $n_f = 5$	0.259	17.7	0.251	18.6	0.261	17.3

Table 2: Values of Mean Square Charge Radii of the Heavy-Light Pseudoscalar D Mesons in  $fm^2$  for finite quark masses  $\langle r^2 \rangle_F$  in the V scheme with  $c = 1GeV$  for  $b = 0.183GeV^2$  with  $\alpha_V = 0.625$  and  $A_0 = 1$  in comparison to the predictions of others.

	$\epsilon_V$	$D_S^+$	$D^0$	$D^+$
$\langle r^2 \rangle_F$ , this work,	0.592	0.077	-0.302	0.172
$\langle r^2 \rangle_F$ [9]	-	0.124	-0.304	0.180
$\langle r^2 \rangle_{T,F}$ (coulombic as parent)[27]	-	0.0101	-0.0139	0.0119
$\langle r^2 \rangle_F$ [28]	-	0.1260	-0.2343	0.1340
$\langle r^2 \rangle_F$ [26]	-	0.1239	-0.3036	0.1849
$\langle r^2 \rangle_F$ (nonrelativistic)[29]	-	-	-0.210	0.114

Table 3: Values of Mean Square Charge Radii of the Heavy-Light Pseudoscalar B Mesons in  $fm^2$  for finite quark masses  $\langle r^2 \rangle_F$  in the V scheme with  $c = 1GeV$  for  $b = 0.183GeV^2$  and  $A_0 = 1$  for both  $\alpha_V = 0.261$  and  $\alpha_V = 0.6$  in comparison to the predictions of others

	$\alpha_V$	$\epsilon_V$	$B_C^+$	$B_S^0$	$B^0$	$B^+$
$\langle r^2 \rangle_F$ , this work	0.261	0.193	0.565	-5.401	-13.651	27.489
$\langle r^2 \rangle_F$ , this work	0.6	0.552	0.018	-0.073	-0.185	0.373
$\langle r^2 \rangle_F$ [9]	-	-	0.0433	-0.119	-0.187	0.378
$\langle r^2 \rangle_F$ [28]	-	-	-	-	-1.464	-
$\langle r^2 \rangle_F$ [26]	-	-	-	-	-0.1866	-
$\langle r^2 \rangle_{T,F}$ (coulombic as parent)[27]	-	-	0.0106	-0.0250	-0.0301	0.0605

It shows that the two loop potential invariably scales up the effective coupling constant from  $\alpha_{\overline{MS}}$  to  $\alpha_V$ . The increase in percentage is also shown in Table 1. It is much more for c-quark than for b-quark. The values of the running coupling constant for B and D mesons are  $\alpha_{\overline{MS}} = 0.22$  and  $\alpha_{\overline{MS}} = 0.39$  respectively [23]. The two loop calculations raise these values for B and D mesons to  $\alpha_V = 0.261$  and  $\alpha_V = 0.625$  respectively as shown in Table 1, and are in good agreement with our results of [11] for D mesons. For B mesons however, the boost given by two loop results is insufficient.

### 3.2 Effect of the Parameter $c$ in the Analysis

In the previous studies on this model [11, 12, 13, 14, 16], the undetermined factor appearing in the series solution of the Schrödinger equation  $A_0$ , (as occurred in equation (5)), was set equal to zero and the effect of  $c$  disappeared altogether in the formalism. In the previous studies, a very small value of the confinement parameter  $b$  could only be accommodated due to the perturbative constraints. However, in the present analysis, the standard quarkonium spectroscopic result [24, 25]  $b = 0.183GeV^2$  can also be accommodated by a suitable choice of  $c$  (taking  $A_0 = 1$ ), since the new perturbative constraints given by equations

(12) and (13) of [15] do not prohibit large values of  $C'(c)$ . As mentioned in [15], since the reduced masses of mesons are about 1 GeV or less, a very large value of the parameter  $c$  occurring in the Hamiltonian given in equation (12) of [11] would most probably not be natural.

### 3.3 Charge Radii

Using equation (24), we calculate the square of their charge radii  $\langle r^2 \rangle_F$  for finite heavy quark masses in  $fm^2$  and compare our results with other results. We show our results for the mean square charge radii of the heavy pseudoscalar mesons in Table 2, taking  $b = 0.183 GeV^2$  and  $c = 1 GeV$  with  $\alpha_V = 0.625$  for the D mesons, and in Table 3 with  $\alpha_V = 0.6$  and  $\alpha_V = 0.261$  for the B mesons. Predictions for D mesons agree well with Hwang [9], but for B mesons, (with  $\alpha_V = 0.261$ ), the corresponding predictions overshoot by a factor of 2 - 3. This anomalous feature can be overcome only if  $\alpha_V$  for B mesons is higher, say  $\alpha_V \sim 0.6$ .

## 4 CONCLUSIONS

We have calculated the charge radii of heavy light flavoured mesons with two loop static potential within a QCD inspired quark model pursued by us in recent years. We have also improved our earlier calculations [11, 13, 14] as substantial confinement effects can now be accommodated in contrast to the previous version of the model. Relativistic effects have been estimated in the wave function at the origin by a procedure analogous to the hydrogen atom in QED. Before conclusion, we also make a few comments.

The approach of the earlier version ( $b \sim 0$ ) could be justified when the two constituent quarks are very heavy and their motion is restricted to the small area around the origin, such as in the case of charmonium or bottomonium. For significant confinement effect ( $b > 0$ ), no such restriction exists and is presumably applicable to the heavy-light mesons as well, as we have done in the present paper. The approach can be further tested for the excited heavy-light mesons and for their mass differences.

As we are considering only pseudo scalar mesons, the spin-spin interactions in equation (11) of [11] which give mass splitting between pseudoscalar and vector mesons is neglected. By redefining the c-term, this can be included.

In the present analysis, our results agree with current available data as well as with those of Hwang[9] for a value of the coupling constant at around  $\alpha=0.6$  to 0.65 instead of the running coupling constants of  $\alpha_{\overline{MS}} = 0.39$  for D mesons and  $\alpha_{\overline{MS}} = 0.22$  for B mesons. Such higher value of strong coupling constant can be generated in the V-scheme with  $O(\alpha_S^3)$  terms for D mesons but falls short for B mesons. As noted earlier [14], this presumably indicates large flavour dependent higher order effects of  $O(\alpha_S^3)$  for B mesons. Moreover, this article gives improved values of heavy-light quark mesons, which could be valuable for future theoretical research.

Although the present analysis is an improvement of the earlier work of [11, 13, 14] in the sense that significant confinement as well as higher order effects in  $\alpha_S$  through V-scheme have been incorporated, it still falls short in explaining correctly the properties of mesons with b-quarks.

## COMPETING INTERESTS

The authors declare that no competing interests exist.

## References

- [1] Brodsky SJ. Tests of quantum chromodynamics in exclusive and inclusive electroproduction. SLAC PUB 5013; 1989.
- [2] Rujula A De, Georgi H, Glashow SL. Hadron masses in gauge theory. Phys. Rev. 1975; D12:147-62.
- [3] Godfrey S, Isgur N. Mesons in a relativized quark model with chromodynamics. Phys. Rev. 1985;D 32:189
- [4] Silverman D, Yao H. Phys. Rev. 1988;D38:214.

- [5] Galkin VO, Mishurov A Yu, Faustov RN. Pseudoscalar decay constants of B and D mesons in the relativistic quark model Sov. J. Nucl. Phys. 1991;53:1026.
- [6] Ji CR, Cotanch SR. Simple relativistic quark model analysis of flavored pseudoscalar mesons. Phys.Rev. 1990;D41:2319-2322. DOI: 10.1103/PhysRevD.41.2319.
- [7] Munczek HJ, Jain P.  $q\bar{q}$  Bound States in the Bethe-Salpeter formalism. Phys. Rev. 1992;D46:438.
- [8] Rosner JL. Determination of pseudoscalar charmed meson decay constants from B meson decays. Phys. Rev. 1990;D42:3732.
- [9] Hwang CW. Charge radii of light and heavy mesons. Eur. Phys. J. 2002;C23:585-590. DOI : 10.1007/s100520200904.
- [10] Rai AK, Parmer RH, Vinodkumar PC. Masses and decay constants of heavy light flavour mesons in a variational scheme J. Phys. G: Nucl. Part. Phys. 2002;28:2275.
- [11] Choudhury DK, Das P, et al. Static and dynamic properties of pseudoscalar mesons. Pramana-J. of Phys. 1995;44:519.
- [12] Choudhury DK, Das P. Static and dynamic properties of heavy light mesons in infinite mass limit. Pramana-J. of Phys. 1996;46:349.
- [13] Choudhury DK, Bordoloi Namita S. The isgurwise function in a QCD inspired quark model. Int. J. Mod. Phys. 2000;A 15: 3667. DOI: 10.1142/S0217751X00002743.
- [14] Choudhury DK, Bordoloi Namita S. The isgurwise function in a QCD inspired quark model with two-loop static potential. Mod. Phys. Lett. 2002;A 17:1909. DOI: 10.1142/S0217732302005510.
- [15] Choudhury DK, Bordoloi Namita S. The isgur wise function and its derivatives in an improved version of the QCD inspired quark model. Mod.Phys. Lett. 2009;A24:443.
- [16] Choudhury DK, Bordoloi Namita S. Form factors and charge radii in a QCD inspired quark model with two-loop static potential. Indian Journal of Physics. 2008;82(6):779.
- [17] Peter M. Static quark-antiquark potential in QCD to three loops; 1997. Available: [aps.org/doi/10.1103/PhysRevLett.78.602](https://arxiv.org/doi/10.1103/PhysRevLett.78.602)
- [18] Peter M. The static potential in QCD-a full Two-Loop Calculation. Nucl. Phys. 1997;B501:471-494. DOI:10.1016/S0550-3213(97)00373-8.
- [19] Schröder Y. The Static potential in QCD to two loops. Schroeder Y Phys. Lett. 1999;B447(3-4):321326. DOI. 10.1016/S0370-2693(99)00010-6.
- [20] Schröder Y. The two-loop static potential. Nucl. Phys. Proc. Suppl. 1999;86:525528. DOI. 10.1016/S0920-5632(00)00616-2.
- [21] Stanley DP, Robson D. Model of meson masses in heavy quark anti-quark systems. Phys. Rev. 1980;D21:3180.
- [22] Stanley DP, Robson D. Relativistic form factors for hadrons with quark-model wave functions. Phys. Rev. 1982;D26:223.
- [23] Groom DE, et al. Particle data group. Eur. Phys. J. C. 2000;15:1.
- [24] Eichten E, et al. Spectrum of charmed quark-Antiquark Bound States. Phys. Rev. Lett. 1975;34:369.
- [25] Eichten E, et al. Charmonium: The model. Phys. Rev. 1978;D17:3090.
- [26] Aitchison IJR, Dudek JJ. Variationally improved perturbation theory and the potential  $\alpha/r + \beta r$ . Eur. J. Phys. 2002;23:605.



- [27] Hazarika BJ, Choudhury DK. Form factors and charge radii in a quantum chromodynamics inspired potential model using variationally improved perturbation theory. *Pramana-J. Physics.* 2015;84:1. DOI:10.1007/s12043-014-0839-x.
- [28] Fernandez FM. Variationally improved perturbation theory for central-field models. *Eur. J. Phys.* 2003;24:289. DOI:10.1088/0143-0807/24/3/309.
- [29] Lombard FM, Mares J. On the meson charge radii. *Phys. Lett.* 2000;B472:150-154. DOI: 10.1016/S0370-2693(99)01395-7.

---

©Pathak et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<http://sciencedomain.org/review-history/10192>