



How does Assisted Natural Regeneration (ANR) Improve Soil Fertility and Plant Cover in the Simiri Commune in Niger?

**Daouda Boukary ^{a*}, Hassimiou Halidou Djabri ^b,
Karimou Harouna Boureima ^b
and Alzouma Mayaki Zoubeirou ^c**

^a Directorate General for the Environment, Water and Forests, National Forest Seed Center of Niger,
Post Box 578 Niamey, Niger.

^b Boubakar Bâ University of Tillabéri, Faculty of Agricultural Sciences, Agroecology Laboratory,
Post Box: 175, Tillabéri, Niger.

^c Abdou Moumouni University of Niamey, Faculty of Science and Technology, Biology Department,
Garba Mounkaila Laboratory, Post Box: 10662 Niamey, Niger.

Authors' contributions

This work was carried out in collaboration among all authors. Author DB designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors HHD and KHB managed the analyses of the study. Author AMZ managed the literature searches, validated the research protocol and coordinated activities throughout the manuscript writing and submission process. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i113205

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107098>

Original Research Article

Received: 27/07/2023

Accepted: 04/10/2023

Published: 12/10/2023

*Corresponding author: E-mail: daoudaboukary@yahoo.fr;

ABSTRACT

Aims: This study aimed to assess Assisted Natural Regeneration (ANR) impacts on improving soil fertility and plant cover through the physicochemical characterization of soil samples taken from ANR fields and control fields.

Study Design: The experimental design is made up of a vegetation observation gradient along an East-West transect and a soil sampling gradient along a North-South transect.

Place and Duration of Study: This study was carried out in the Sahelian zone in the rural commune of Simiri in Niger between May and June 2023.

Methodology: Data on the diversity of species predominant in the ANR system, the social categories of ANR users and farmers' perceptions of ANR were collected on a sample of 120 farmers chosen at random from three villages in the commune of Simiri.

Results: A sandy texture was recorded in all the samples submitted for analysis, with an overall acid of soil pH (5.18 - 5.62). The proportions of the main nutrients: C(0,16), N(0,015), P(6,88), K(0,066), Mo(0,30) C/N(10,83) were relatively higher in the fields under ANR practices compared with the control fields. With regard to the dominant woody species, the results show that the stands are mainly based on *Combretum glutinosum*, *Combretum micranthum* and *Guera senegalensis*. The ability of woody species to withstand stressful conditions could act as a bulwark against climatic hazards. The strong dominance of these species is due, among other things, to their regenerative capacity, their rapid spread and, above all, their economic interest in the production of firewood and other ecosystem services.

Conclusion: The Assisted Natural Regeneration (ANR) adoption in farming practices has often been accompanied by the creation of village committees for the protection and management of trees and other areas of sustainable natural resource management. Despite mutilation and other illegal felling, the new agroforestry parks increase the resilience of agricultural production systems to climatic shocks and biotic stresses.

Keywords: Woody species; ANR; stand; plant cover; dissemination; Simiri.

1. INTRODUCTION

The disruption of the ecological balance in the Sahel is mainly the result of the recurrent droughts that characterized the climate between 1970 and 1980 [1]. The consequences of this upheaval included energy and food crises, which caused communities to turn to forest areas to stem the energy crisis and meet the basic needs of an ever-growing population [2]. As a result, the natural resources that contribute to the livelihoods of this population are in a state of advanced degradation under the combined effects of human actions and climate change [3]. In addition, strong demographic pressure on land such as uncontrolled occupation of arable land, irrational exploitation of natural resources, increasing deforestation has resulted in a sharp reduction in vegetation cover throughout the Sahel [4]. The scarcity or even disappearance of plant cover is causing the desert to advance, reducing soil fertility and lowering agricultural productivity, leading to food insecurity. In addition to this ongoing degradation of natural resources, the major challenge in developing countries remains, to a large extent, the issues of poverty and food security [3]. The generally slow

performance of natural ("passive") regeneration in fern-infested areas in the tropics [5,6] has sparked the testing of a range of alternative management techniques to accelerate regeneration processes [7].

Integrating multipurpose woody crops into the agricultural production system helps to mitigate deforestation, land depletion and poverty among rural populations [8]. Faced with environmental degradation and strong land pressure, farmers in densely populated areas, particularly in south-central Niger, have intensified their agricultural production systems. They have done so by increasing the number of trees and shrubs in their fields, thus creating new agroforestry parks, the scale of which in the Zinder, Maradi and Tahoua regions is around 5 million hectares [9].

The approach of assisted natural regeneration (ANR) aims to strike a balance between high-cost restoration planting to restore biodiversity to small areas and the establishment of commercial plantations over large areas to restore productivity [10]. However, despite the large-scale dissemination of these technologies, their adoption by certain communities remains timid

[11]. Numerous studies show that natural regeneration managed by farmers has increased crop yields from 31 to 350 kg/ha and ensured food security for families, even during drought years. However, cereal yields remain low and will not be enough to feed a rapidly growing population (Abasse et al., 2023). [12] Moreover, ANR increases the incomes of all social categories, even the most vulnerable (men, women and young people). Pruning trees in the fields has also reduced the distances travelled by women to collect firewood by increasing the number of trees in the fields on the outskirts of the villages., Thereby improving the availability of firewood in the surrounding fields. ANR has also increased the availability of fodder for farmers and agro-pastoralists, with ANR households harvesting 30 to 45 kg of fodder per day [12].

Increasing the number of trees and shrubs per hectare has increased litter production. This improves soil structure and allows greater quantities of water to be stored. The addition of litter also helps to improve soil fertility. Certain species, which often dominate regeneration, such as *Piliostigma reticulatum*, *Guiera senegalensis* and *Combretum glutinosum*, have a positive impact on the content of chemical elements (carbon, nitrogen and available phosphorus) in the soil. ANR has enabled village communities to adapt better to climate change and to strengthen their resilience. Indeed, even when harvests fail, farmers can cut down certain trees and sell them on the market as firewood or service wood, enabling them to buy cereals. ANR also has a positive impact on crop yields, even in years with low rainfall [12]. Hence the interest of this study, which aims to assess the impact of ANR on improving soil fertility and increasing the density of woody plants in crop fields.

2. MATERIALS AND METHODS

2.1 Study Area

The study was carried out in the rural commune of Simiri and covered three villages in the area: Guessé Gao banda (1°1'21" N; 2°1'5.4" E), Guessé Sinsan (1°1'11" N; 2°1'15" E) and Simiri (1°8'18" N; 2°8'25" E)

The rural commune of Simiri is 80 km north of Niamey in the Ouallam department. The climate is arid, with annual rainfall varying between 400 and 600mm [13]. The rainfall is irregularly distributed in time and space. The rainy season extends from June to September. The average maximum temperature varies from 24°C in

January to 33°C in April. The representative toposequence of the region is characterised by a succession of four major zones typical of the tropical ferruginous plateaux of the Sahel: the armoured plateau, the sandy skirt, the glaciais and the lowlands [14]. Skeletal soils of the rigosol and lithosol types formed on terminal continental clayey sandstones are found on the plateaux [15]. The sandy skirt and glaciais contain sandy tropical ferruginous soils with little leaching, followed by alluvial soils in the lowlands. The natural vegetation consists of shrub steppe on the glaciais and shrubs to trees in the lowlands, while a contracted formation subject to severe degradation can be seen on the plateaux [16]. The main woody species are *Guiera senegalensis*, *Combretum micranthum* and *Combretum glutinosum*. The herbaceous layer is dominated by *Mitracarpus scaber*, *Eragrostis tremula* and *Cenchrus biflorus*. Fig. 1. below shows the geographical location of the Simiri commune.

2.2. Documentary Research

This phase involved gathering the documentation needed to understand properly the study subject and to establish the state of knowledge on the subject. It was carried out using databases (website, scientific journals), activity reports and scientific publications.

2.3 Field Investigation and Demographical Analysis

To collect both quantitative and qualitative data, individual interviews were conducted with a sample of 120 farmers practising ANR in their fields. These farmers were located in three villages in the rural commune of Simiri: Guessé Sinsan, Guessé Gaobanda and Simiri. The interviews were conducted using a semi-structured questionnaire. A systematic inventory of woody species was carried out in the fields in order to assess the diversity of species used in ANR.

The demographic analysis (DA) made it possible to collect and determine the general characteristics of the target groups concerned by the survey. Once the data had been analyzed, qualitative and quantitative information led to the grouping of respondents on the basis of a number of parameters, including age, gender, level of education and experience in the practice of natural assisted regeneration. The results of the demographic analysis were presented in graph form.




Fig. 1. Location map of the rural commune of Simiri
(Source: Moussa Boubacar, 2019)



Fig. 2. Interview with an ANR practitioner in guessé sinsan (Daouda, 2023)

2.4 Soil Sampling and Analysis

A total of twelve (12) soil samples were taken from all three sites, with three samples per site and three controls. The soil samples collected were analysed at the INRAN Niamey soil science laboratory. The physico-chemical parameters used to characterize the soil were :

granulometry by the Robinson pipette method; pH by determination of hydrogen ions; cation exchange capacity (CEC) and the sum of exchangeable bases (S) by the silver thio urea method; total nitrogen by the Kjeldahl method; organic carbon by the Walkley & Black method; assimilable and total phosphorus (P) by the Bray1 method, electrical conductivity by the

conductimetry method, exchange acidity, organic matter and exchangeable bases. The Fig. 3 and 4 below show respectively the packed soil samples and an overview of a field under ANR practices.

2.5 Data Analysis and Processing

The survey data were tabulated and analysed using Excel, then subjected to descriptive analysis using SPSS software. The soil analysis results were also subjected to Linear discriminant analysis in order to identify similarities between variables at different sites.

3. RESULTS

3.1 Breakdown of Producers by Sex and Age

Fig. 5 below shows the breakdown of NAR users by gender and age group.

This graph shows that the average age of respondents is 39.97 ± 12.76 , with a minimum of 20 and a maximum of 67. Regardless of the sites, women are in the minority in the practice of ANR.



Fig. 3. packed soil samples (Daouda, 2023)



Fig. 4. Overview of a field of RNA (Daouda, 2023)




Fig. 5. Breakdown of respondents by gender and age group




Fig. 6. Dispersion of cationic exchange capacity (CEC) and exchange acidity (EA) values by site (Guessee_S= Guesse Sinsan x-axis; Guessee= Guesse Gao Banda x-axis)

3.2 Dispersion of soil physicochemical parameters by site

This figure shows that the cationic exchange capacity values recorded at the Simiri and Guesé Gao-Banda sites are scattered around the average of CEC values for samples from the Simiri and Guesse Gao banda sites, unlike those obtained at the Guesé Sinsan site, which are close to the general CEC average value for all

sites. On the other hand, the variation in exchange acidity values shows that the Simiri and Guesé Gao-Banda sites have values well above the general average of the exchange acidity value, while the Guesé Sinsan site has values below the average of the same variable. This indicates a fairly clear change of the exchange acidity between sites. In addition, low cationic exchange capacity and exchange acidity values were observed on control samples taken

outside fields under assisted natural regeneration.

3.3 Constraints to Adoption of ANR in the Farming Environment

The figure below illustrates the main constraints to adoption of ANR according to the farmers surveyed.

The figure shows that conflicts over wood harvesting, the problem of defending fields under

ANR and poor knowledge of young shoot maintenance techniques are the main constraints faced by farmers in adopting ANR, with proportions of 32.5%, 31.25% and 16.25% respectively.

3.4 Socio-Economic Impacts of ANR

3.4.1 Impacts on household living conditions

Fig. 8 below presents the impacts of the ANR on household living conditions.




Fig. 7. Constraints to adoption of ANR




Fig. 8. Impact of the NAR on household living conditions




Fig. 9. Main advantages of ANR

The graph shows that 68.25% of respondents said that the use of ANR in crop fields had increased their food stocks, compared with 31.25% who said that it had clearly increased their income, in particular through the sale of firewood, service wood and non-timber forest products. This income is then used to reconstitute part of the food stock in addition to that derived from the production of fields whose fertility is improved by these practices.

3.4.2 Impacts on soil and climate components

The figure below illustrates the main advantages of ANR.

There are several advantages to implementing ANR in cropping systems. Indeed, the following main benefits emerge from this graph: Return of wild fauna, improved soil fertility, and firewood production with a preponderance of 30%, 27.5% and 20% respectively.

3.4.3 Impacts of ANR on soil physicochemical properties

The soil analysis results show that the fields under RNA have fairly variable characteristics and an acceptable level of fertility. The pH was generally acidic in all samples (5.07 - 5.62), with relatively levels of organic matter (0.10 - 0.44), carbon (0.06 - 0.28) and total nitrogen (0.006 - 0.025). In addition, the predominantly sandy texture of the soil samples justifies the low CEC

values, which vary according to OM content. The slider between calcium and magnesium depends on the size of the CEC. Fig. 10 below illustrates the variation in the cation exchange capacity of the soil in fields under ANR according to the Ca^{++}/Mg^{++} slider.

This graph shows that the CEC of the soils varies from one site to another, unlike the Ca^{++}/Mg^{++} complex, which shows little variation from one site to another. In terms of soil texture, the results show that all the samples are predominantly sandy, with relatively low clay and silt content. So the soils texture is highly dominated by fine and coarse sands proportions. The figure shows two peaks (convex and concave) representing the two extreme values for the Ca^{++}/Mg^{++} complex and the CEC for all the samples analysed.

3.4.4 Diversity of woody species in agroforestry parks under ANR

The figure below shows the diversity of woody species in fields under ANR practices in the Simiri zone.

This figure shows that several woody plant species are used in ANR systems. These include *Combretum micranthum* and *glutinosum*, *Guera senegalensis* and *Annona senegalensis*, which are the most predominant species in agroforestry parks in the Simiri zone.

Table 1. Physico-chemical characteristics of the soil in the fields under practice

N° Lab.	Identification	Prof. en cm	pH	Mèq/100 g						Phosphore ppm			Fer %		%			
				H ₂ O	C.E m ³ /cm	CA ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺	Tot	CEC	AE	Assim.	Tot	Libr	Tot	Carb.	M.O
2023/188	Simiri Témoin	0-25	5.18	0.01	1.10	3.48	0.165	0.071		4.86	0.04	8.11	0.122		0.26	0.44	0.022	12
189	Smiri 1	0-25	5.62	0.04	1.25	4.11	0.174	0.092		5.69	0.06	6.31	0.110		0.10	0.18	0.009	11
190	Simiri 2	0-25	5.45	0.04	1.15	3.56	0.174	0.071		5.03	0.08	5.80	0.122		0.28	0.47	0.024	12
191	Guessé Gao banda Témoin	0-25	5.46	0.04	1.20	3.34	0.157	0.071		4.83	0.06	2.56	0.134		0.21	0.36	0.018	12
192	Guessé Gao banda 1	0-25	5.45	0.05	1.05	1.21	0.044	0.021		2.37	0.04	14.51	0.108		0.18	0.31	0.016	11
193	Guessé Gao banda 2	0-25	5.51	0.06	1.25	2.03	0.078	0.036		3.49	0.10	4.61	0.125		0.28	0.49	0.025	11
194	Guessé Gao banda 3	0-25	5.38	0.05	1.25	2.89	0.165	0.046		4.41	0.06	8.28	0.106		0.09	0.45	0.023	04
195	Guessé Gao banda	0-25	5.38	0.05	1.25	3.16	0.183	0.077		4.79	0.12	2.99	0.098		0.07	0.12	0.006	12
196	Guessé Sissan Témoin	0-25	5.46	0.05	1.50	3.43	0.183	0.066		5.24	0.06	2.82	0.125		0.08	0.13	0.007	11
197	GuesséSissan 1	0-25	5.10	0.02	1.50	3.77	0.165	0.092		5.63	0.10	12.38	0.153		0.06	0.10	0.005	12
198	Guessé Sissan 2	0-25	5.07	0.02	1.40	2.98	0.087	0.092		4.60	0.04	7.43	0.129		0.10	0.18	0.009	11
199	Guessé Sissan 3	0-25	5.33	0.05	1.40	3.12	0.174	0.066		4.86	0.10	6.83	0.108		0.22	0.39	0.020	11

EC: electrical conductivity; CA⁺⁺: calcium; Mg⁺⁺: magnesium; Na⁺: sodium; K⁺: potassium; CEC: cation exchange capacity; AE: exchange acidity; assimilable phosphorus; carb: carbon; Mo: organic matter; C/N: carbon to nitrogen.




Fig. 10. Ca⁺⁺/Mg⁺⁺ complex and CEC in soil




Fig. 11. Diversity of woody species used in ANR

5. DISCUSSION

Increased degradation of cultivated land and other special ecosystems has led to massive investment in restoration activities in Niger. For this reason, the country has signed up to the African Forest Landscape Restoration (AFR100) initiative, which aims to restore 100 million hectares by 2030. To honour this commitment, there are plans to further promote agroforestry practices, including assisted natural regeneration for the rapid reforestation of agricultural areas, which will ensure the productivity of Niger's agro-

ecosystems. In this study, the results showed a good perception of ANR by the producers surveyed, hence the involvement of all socio-professional strata in the adoption of this practice. The farmers surveyed said that they had intensified their agricultural production system by increasing the number of trees in their fields. This increase does not result from new plantings but rather from the protection and maintenance of young tree shoots that appear spontaneously in the fields, contributing in the long run to the formation of agroforestry parks with, in particular, various centers of interest for

the communities and the landscape. These results are in line with those of several authors (Sitou *et al.*, 2018; Ado *et al.*, 2019; Botoni *et al.*, [4] who find that ANR is a good way of improving agricultural production and combating food insecurity. Furthermore, Sitou *et al.*, 2018; Ado *et al.*, 2019; Botoni *et al.*, 2010; state that ANR could be a strategy for combating food insecurity through the sale and marketing of woody forest products. Various woody species are protected and maintained in crop fields, providing multiple ecosystem services, including soil fertilization through litter from dead leaves, climate regulation, carbon sequestration, recreation and runoff reduction. These results corroborate those of Bagnian *et al.* [17] Moussa *et al.* [18] Mamane [19] and Maâzou [20] who state that woody stands in fields under RNA are most often dominated by *Combretum glutinosum*, *Combretum micranthum*, *Piliostigma reticulatum*, *Guiera senegalensis*, *Faidherbia albida* and *Prosopis africana*. According to Bagnian *et al.* [17] Moussa *et al.* [18] Mamane [19] and Maâzou [20] the specific contribution of each species is highly variable, ranging up to more than 50% of the total tree stand. Improving soil fertility under RNA is one of the primary objectives of natural assisted regeneration. Farmers invest in trees mainly to maintain or improve the fertility of their soils, and the impact of certain species on chemical fertility has been confirmed by several studies (Mansour *et al.*, [21] Camara *et al.*, 2017; Dan Lamso *et al.* [22,23] Zounon *et al.*, [24] Traoré, 2012; Bodo *et al.*, [25] Soil analysis results showed that the content of certain nutrients was much higher in fields under RNA compared with control areas not under RNA. These include C (%), N (%), K (meq/100g) and C/N. This suggests that an increase in tree density is always followed by an improvement in soil fertility. A study by Dan Lamso *et al.* [26] showed that ANR is an effective fertility management practice for tropical ferruginous soils cultivated in Niger. In the same way agroforestry being an agro-ecological, climate-smart, environmentally friendly, and sustainable practice, has many ecosystem services, one of which is soil fertility enhancement [27].

The results of the household surveys showed that the practice of ANR in the fields generates additional income such as improving soil fertility, creation of a microclimate, availability of non-timber forest products for practising households in addition to the increase in cereal production in the fields. This increased income comes from the

sale of wood and non-wood forest products from the ANR. Although this income has not been accurately evaluated in this study, the producers claim that the sale of ANR products considerably reduces the poverty level of households and, in turn, enables them to obtain food supplies. This protects them from food insecurity and increases the duration of their food stocks. These results are in line with those of Bagnian (2010), who states that the action of trees increases the duration of agricultural production stock by 5 to 7 months on average, which contributes significantly to food stability.

The management of ANR is often accompanied by better consultation and organisation within villages in order to manage the new tree capital effectively.

Several constraints limit the use of ANR in the villages covered by the study. These are mainly conflicts over wood harvesting (33%), problems with defending fields under ANR (31%) and lack of knowledge of techniques for maintaining young shoots (16%). These results are similar to those of Kabirou (2022), who notes that transhumant herders, who generally come from elsewhere during the dry season, carry out severe pruning, which more often than not compromises the survival of the trees. Similarly, the animals of local livestock farmers, which are sometimes left to roam, graze on the new shoots left in the fields. This leads to frequent physical conflicts. In addition, young people and women from the villages fraudulently cut down plant species in the fields without considering the consequences. Also, woodcutters from several localities cut down trees clandestinely at ground level in the absence of the field owner, often late at night (Kabirou, 2022). Finally, the partial application of the provisions of the forestry code relating to illegal felling of trees, irrational and uncontrolled exploitation of forest products by rural actors (customary authorities and village forest resource management committee) and the inadequacy of forestry regulations that take into account the status of regenerated trees in the fields are constraining ANR in the study area (Kabirou, 2022). The introduction of ANR has often been accompanied by the creation of village organisations for the protection and management of trees. These organisations have adopted community actions against tree felling and theft, and this new social capital helps to ensure the sustainability of the tree capital established in the fields. The new agroforestry parks increase the resistance of agricultural

production systems to climatic shocks and sequester large quantities of carbon [28].

6. CONCLUSION

At the end of this study, the results show that ANR is an effective agroforestry practice for integrated management of soil fertility, crop productivity and the improvement of farmers' sustainable livelihoods. Given its multiple benefits, it encourages a clear awareness and willingness on the part of small-scale farmers to protect and manage woody species. The adoption of this practice is closely linked to the farmers' perception of the causal link between the presence of woody species in high densities in the fields and the level of fertility of the fields, and between the level of fertility of the soil and the productivity of the land. Furthermore, the soil and climate conditions (sandy soil, semi-arid climate) are undeniable determinants for the adoption of ANR in the villages of Guessé and Simiri. The fact that this practice is accessible to all social categories is conducive to its widespread adoption. In addition, farmers derive income from ANR through the sale of firewood and other tree products. Women benefit from ANR because it increases the availability of fuel, reducing the distances they have to travel to collect wood. It also increases non-timber forest products, which help to improve family nutrition and the quality of life.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. RINAUDO T.: Une brève histoire de la régénération naturelle assistée, l'expérience du Niger. *Echo, Note technique*. 2010 ;27.
2. Dramé YA, Berti F. 2008. Les enjeux socioéconomiques autour de l'Agroforesterie villageoise à Aguié (Niger), *TROPICULTURA*, 2008, 26, 3, 141- 149. ISSN 0771-3312 eISSN 2295-8010Dramé YA, Berti F. 2008. Les enjeux socioéconomiques autour de l'Agroforesterie villageoise à Aguié (Niger), *TROPICULTURA*. 2008 ; 26(3):141- 149. ISSN 0771-3312 eISSN 2295-8010 ;
3. LAWALI Sitou, Abdoulaye DIOU, Boubé MOROU, Kassimou ABDOU KONA, Laminou SAIDOU, Chaibou GUERO et Ali MAHAMANE, Régénération Naturelle Assistée (RNA) : outil d'adaptation et résilience des ménages ruraux d'Aguié au Niger. *Int. J. Biol. Chem. Sci.* 2018 ; 12(1) :75-89 ;
4. Botoni E, Larwanou M, Reij C. La régénération naturelle assistée (RNA) : une opportunité pour reverdir le Sahel et réduire la vulnérabilité des populations rurales. Le projet majeur africain de la Grande Muraille Verte : Concepts et mise en œuvre [en ligne]. Marseille : IRD Éditions, 2010. DOI : 10.4000/books. Ird editions. 2010 ;2122 .
5. Shono K, Cadaweng EA, Durst PB. Application of assisted natural regeneration to restore degraded tropical forestlands. *Restoration Ecology*. 2007;15(4):620-6.
6. Crouzeilles R, Ferreira MS, Chazdon RL, Lindenmayer DB, Sansevero JBB, Monteiro L, Iribarrem A, Latawiec AE, Strassburg BBN. Ecological restoration success is higher for natural regeneration than for active restoration in tropical forests. *Science Advances*. 2017 ;3(11) : [e1701345]. Available : <https://doi.org/10.1126/sciadv.1701345>;
7. Rurangwa L, Matthews T, Niyigaba P, Tobias J, Whittaker RJ. 'Assessing tropical forest restoration after fire using birds as indicators: an afro-tropical case study', *Forest Ecology and Management*. 2021;483:118765. Available:<https://doi.org/10.1016/j.foreco.2020.118765>
8. Bonkougou Edouard. Introduction à l'agroforesterie : stage de formation ; l'agroforesterie pour une production agricole durable et une meilleure gestion des ressources naturelles au Sahel, Ouagadougou, Bobo-Dioulasso du 9 au 27 novembre ; 1998.
9. Cotillon S, Tappan G, Reij C. Land use change and climate smart agriculture in the Sahel. In: Villalon, L.A. (ed.) *The Oxford Handbook of the African Sahel*, Oxford University Press. 2021;209-230 ;
10. Shono K, Cadaweng EA, Durst PB. Application of assisted natural regeneration to restore 815 degraded tropical forestlands. *Restoration Ecology*. 2007;15 :620-626. DOI:10.1111/j.1526- 816 100X.2007.00274.x

11. Zarafi Marou A, Abasse AT, Bokar M, Niang A et Cheik O. Traoré, 2002. Analyse de l'adoption de la régénération naturelle assistée dans la région de Maradi au Niger. 2^{ème} Atelier régional sur les impacts socio-économiques de l'agroforesterie au sahel. Bamako- mali 4 au 6 mars. 2002;7.
12. Abasse T, Massaoudou M, Rabiou H, Idrissa S, Dan Guimbo I, Régénération naturelle assistée au Niger: l'état des connaissances. Tropenbos International, Ede, Pays Bas. 2023. DOI: 10.55515/GHFR2702 ;
13. Saadou M. La végétation des milieux drainés nigériens à l'Est du fleuve Niger. Thèse de Docteur ès Sciences Naturelles, Université de Niamey, Niger. 1990 ;395.
14. Ambouta JMK, Valentin C, Laverdière MR. Jachère et croûtes d'érosion au sahel. Sécheresse .1996 ;7:269-75 ;
15. Ambouta JMK. Définition et caractérisation des structures de végétation contractée au sahel: cas de la brousse tigrée de l'ouest nigérien. In Fonctionnement et Gestion des Ecosystèmes Forestiers Contractés Sahéliens, d'Herbès JM, Ambouta JMK, Peltier R (éds). John Libbey Eurotext : Paris .1997 ;41-57 .
16. Moussa Mamoudou BOUBACAR, Maman Maârouhi INOUSSA, Jean-Marie Karimou AMBOUTA, Ali MAHAMANE, Aagaard Axelsen JORGEN, Yahou HARISSOU et Habou RABIOU, Caractérisation de la végétation ligneuse et des organisations pelliculaires de surface des agroécosystèmes à différents stades de dégradation de la Commune rurale de Simiri (Niger). *Int. J. Biol. Chem. Sci.* 2013 ; 7(5):1963-1975.
17. Bagnian IT. Adam MM. Adamou, I. Chaibou, A. Mahamane. Structure et dynamique de la végétation ligneuse juvénile issue de la régénération naturelle assistée (RNA) dans le Centre- Sud du Niger. *International Journal of Biological and Chemical Sciences.* 2014 ;8(2):649-665.
18. Moussa ML. Mahamane M. Saadou.. Caractérisation des peuplements ligneux des parcs à *Faidherbia albida* (Del) A. Chev. et à *Prosopis africana* (Guill., Perrot et Rich.) Taub. du centre-sud Nigérien. *Journal of Applied Biosciences* 2015;94 :8890–8906 ;
19. Mamane A. Dynamique des peuplements ligneux dans les agrosystèmes au centre Sud du Niger : cas des régions de Maradi, Tahoua et Zinder. Masters thesis, Université Dan Dicko Dankoulodo, Maradi, Niger. 2017 ;66.
20. Maazou MH. Caractérisation de la végétation ligneuse issue de la régénération naturelle assistée (RNA) et Évaluation de son effet combiné aux fertilisants sur la culture associée mil-niébé dans la commune de Guidan Sori. Masters thesis, Université de Diffa, Niger. 2019 ;76.
21. Mansour AM, AM. Zoubeirou A. Kadri J-M.K. Ambouta, N. Dan Lamso. Effet de l'arbre *Acacia senegal* sur la fertilité des sols de gommerais au Niger. *International Journal of Biological Chemical Sciences.* 2013 ;7(6):2328-2387.
22. Dan Lamso NY. Guero A. Tankari Dan-Badjo L. Rabah, B.B. Andre, D. Patrice, A.D. Tidjani, N. Ado Maman, JM.K.Ambouta. Effet des touffes de *Guiera senegalensis* sur la production du mil dans la région de Maradi (Niger). *Revue des BioRessources* . 2015a.5(2) :1-13 ;
23. Dan Lamso N, Y. Guero, A. Tankari Dan-Badjo, L. Rabah, B. B. Andre, D. Patrice, A.D. Tidjani, N. Ado Maman, JM. K. Ambouta. Effet des touffes de *Hyphaene thebaica* (Mart) sur la production du mil dans la région de Maradi (Niger). *International Journal of Biological Chemical Sciences* 2015b ;9(5) :2477.
24. Zounon FCS, T. Abasse, M. Moussa, H. Rabiou, B. Vincent, D. Tidjani, A. Karimou. Effet de la combinaison régénération naturelle assistée (RNA) et microdose d'engrais sur la production du mil (*Pennisetum glaucum* (L.) R. Br) dans les zones agroécologiques du centre-sud du Niger. *European Scientific Journal.* 2020 ;16(6) :317-338;
25. Bodo Seyni, B, X. Morvan, O. Malam Issa, D. Tidjani Adamou, JM. Ambouta Karimou, B. Marin, M. Ponthieu, G. Fronteau. Connaissance locale de la variabilité de surface du sol et des contraintes associées pour la production du niébé en zone sahélienne du Niger. *Etude et Gestion des Sols.* 2019 ;26:65-7 ;
26. Dan Lamso N, A.M.Nassirou G.Yadji. Assisted Natural Regeneration (RNA): an efficient practice for soil fertility management of cultivated tropical ferruginous soils in Niger. *International Journal of Innovation and Applied Studies.* 2022;36(2):609-618.
27. Tsufac AR, Awazi NP, Yerima BPK. Assessing the determinants of soil fertility

- in cocoa-based (*Theobroma cacao*) agroforestry systems in the Littoral region of Cameroon. *International Journal of Plant & Soil Science*. 2020;32(17):52–60.
Available: <https://doi.org/10.9734/ijpss/2020/V32i1730386>
28. Abdou MM, Z. Alzouma Mayaki A. Kadri, J.-MK. Ambouta, J.-M. Karimou, N. Dan Lamso. Effet de l'arbre *Acacia senegal* sur la fertilité des sols de gommériaies au Niger. *International Journal of Biological Chemical Sciences*. 2013;7(6): 2328-2337.

© 2023 Boukary et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107098>