



Impact of Host Age, Gestational Week and Parity on Susceptibility to Urinary Tract Pathogens

Zekeri Sule ^a, Mitsan Olley ^{b*}, Maureen Uchechukwu Okwu ^c,
Clement Osaiyuwu ^b and Maiza Mariam Omoshoke ^b

^a *Department of Obstetrics and Gynaecology, College of Health Science, Igbinedion University Okada, Nigeria.*

^b *Department of Medical Laboratory Science, College of Basic Medical science, Igbinedion University Okada, Nigeria.*

^c *Department of Microbiology, Natural and Applied Science, Igbinedion University Okada, Nigeria.*

Author's contribution

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95686>

Original Research Article

Received: 21/10/2022

Accepted: 29/12/2022

Published: 10/01/2023

ABSTRACT

Background: Urinary tract infections are inflammatory responses of the urothelium to non-flora microorganisms. Aim: This study aimed at identifying the urinary tract pathogens in association with other risk factors in pregnant women residing in the rural community of Okada, Edo State.

Methods: The study entails a cross sectional study involving pregnant women at various gestational age, a mid stream, non-repetitive urine samples were collected in sterile bottles and cultured on bacteriological media at 37°C overnight, isolates were identified using standard conventional procedures.

*Corresponding author: E-mail: olley.mitsan@iuokada.edu.ng;

Results: The organisms isolated include *Staphylococcus aureus*, *Escherichia coli*, *Candida albicans* with prevalence of 35.7%, 5.1% and 27.1% respectively, the age range 26-31 have a prevalence of 37.7% with a p-value of 0.006, the distribution of UTI in relation to other risk factors such as parity, haemoglobin genotype and level of education had p-value of 0.179, 0.067 and 0.0001 respectively.

Conclusion: The relationship between UTI is not statistically significant with respect to age and parity, but statistically significant with gestational age and level of education.

Keywords: pregnant women; urinary tract pathogens; microorganisms; pyelonephritis.

1. INTRODUCTION

“Urinary tract infections (UTIs) are inflammatory responses of the urothelium to bacterial invasion by non-flora microorganisms. It is one of the most common medical issues associated with pregnancy and may involve the reproductive organs if not well managed” (Brook *et al.*, 2001). Despite major progress in the management of urinary tract infections in pregnancy, they are still linked to negative foeto-maternal outcomes [1]. According to Ali and Abdallah, [2], “urinary tract infection describes microbial colonization or inflammation of the bladder (cystitis), urethra (urethritis), or renal pelvis and kidneys (pyelonephritis) with growth of up to 10^6 organisms per μ l of urine in an asymptomatic presentation” (Emilie *et al.*, 2011). The presence and growth of these microorganisms in the urinary tract can lead to these infections, irrespective of age and mostly in pregnancy in association with other risk factors can be life threatening [3].

Culturing of urine specimens is the gold standard for diagnosing urinary pathogen, as well as allowing for quantitative bacteriuria quantification [1]. According to Nwachukwu *et al* [4], “the common etiologic agents of UTI are *Escherichia coli*, *Klebsiella spp*, *Staphylococcus aureus*, *Pseudomonas spp*, and *Streptococcus*”. “These organisms can be found in the external genitalia, vagina, rectum, genital tract, and gastrointestinal tract and can infect cells in the urinary tract by attachment and eventual ascension, causing bacteremia, due to the renal cells being compromised. Advanced age, number of parturitions, number of coitus per week, recessive sickle cell anemia, and diabetes, previous history of UTI, immunodeficiency, and urinary tract abnormalities are conditions that can increase the risk of UTI in pregnant women” [5].

“Pregnancy changes including anatomical, physiological and hormonal increases susceptibility to the development of UTI. These

changes can be due to increased levels of progesterone and estrogen but also due to pressure created by the growing uterus” [6]. About 70% of women during pregnancy develop glycosuria, which supports bacterial growth in the urine (Kant *et al.*, 2017). UTIs are may also be associated with premature delivery, low-birth-weight infants, cesarean delivery, morphological abnormalities, and infant mortality [7]. Early diagnosis, administering of adequate treatment using antibiotics has been shown to helped reduced these risks by 80% (Nicolle, 2015). In this rural community there is paucity of UTI in relation to other associated risk factors such gestational age, parity and Haemoglobinopathy. This study aimed at identifying the urinary tract pathogens in association with other risk factors in pregnant women residing in the rural community of Okada, Edo State, Nigeria.

2. MATERIALS AND METHODS

2.1 The Study Area

The study was carried out in Igbinedion University Teaching Hospital, Okada in Edo State using the facilities of the Medical Microbiology Laboratory.

2.2 Study Design

This is a cross sectional study involving pregnant women visiting the antenatal clinic and general outpatient clinic. The study comprised of 150 women who were pregnant at various gestational age, a mid- stream non repetitive urine was collected from each participant following administration of questionnaire after informed written consent. Urine samples were collected for urine microscopy, culture and sensitivity. Furthermore, those currently on antibiotics were excluded from the study.

2.3 Screening of Urine Bacteria

Culture method described by Chesbrough, [8] was used for the isolation of bacteria.

Briefly, mid-stream urine was cultured on Cysteine lactose electrolyte deficient agar (CLED) and Sabouraud dextrose agar. At 37°C, all cultures were incubated aerobically for 18-24hrs.

2.4 Identification of Isolates

“All bacterial isolates were subjected to taxonomic analysis by the determination of their morphological, physiological and biochemical characteristics as adopted” by Fawole and Oso [9] Holt et al. [10] Cowan and Steel, [11] Cheesbrough, [8].

3. RESULTS

150 urine samples were collected from pregnant women attending Igbinedion University Teaching Hospital (IUTH). The age ranges of the subjects were between 20-47 years as shown in Table 1. The age ranges 20-25, 26-31, 32-37 and 38-43 had 17.1%, 37.7%, 32% and 15.4% prevalence for urinary tract infections respectively and is statistically significant with P- value 0.006. Table 2 shows the distribution of urinary tract infections in pregnant women in relation to gestational age, the week, (1- 12), (13- 25) and (26-40) had 13.6%, 17.9% and 56.8% prevalence rate of infection and are statistically significant with p-value 0.0001. The distribution of UTI in relation to other factors such as marital status, level of education, past history of infection, sexual activity and haemoglobin genotype is shown in Table 3; the level of Education with respect to UTI is statistically significant with p-value of 0.0001. The distribution of UTI in pregnancy in relation to parity is shown in Table 4, parity of 0, 1, 2, 3, and > 4 had UTI prevalence of 31.3%, 21.1%, 15.4%, 22.2% and 43.8% respectively and is not statistically significant with respect to UTI.

Table 5 shows the microbial distribution; *Staphylococcus aureus*, *Candida albicans*, and *Escherichia coli* had occurrence rate of 35.7%, 27.1% and 5.1% respectively.

4. DISCUSSION

Infection of the urinary tract (UTI) represents the most common medical complication of pregnancy especially in asymptomatic bacteriuria. Pregnant women are at greater risk of UTIs, particularly because of the physiologic,

hormonal and anatomic changes that occur in normal pregnancy.

In this research, about 150 urine samples of pregnant women were collected of which the overall prevalence of UTI among pregnant women in this study was 43(28.7%). This is less than the prevalence reported in Saudi Arabia 53.5% [12], but within the range obtained in Ambo Central Ethiopia 18.7%, Ismailia, Egypt 29%, Benin city, Nigeria 21% and in Nepal 37.8% respectively [13]. These variations in prevalence might be as a result of differences in social habits, religious affiliation, and personal hygiene of participants. However, spirituality has a way of making people play down on personal and environmental hygiene.

The age range in this research with high prevalence was between 26-31years with 26(37.7%) which agrees with a study carried out in Onitsha, Nigeria [4]. This may be as a result of sexual activity which increases the risk of UTI and the women of such age group are mostly sexually active.

“With respect to parity; Parity greater than 4 has the highest prevalence with 4(43.8%). however, in a study conducted by in Gorgan, the highest rate of infections was after the third pregnancies (4.73%)” [14]. “The results from the studies of Mobbasheri et al. were consistent with the results from the study in America” [15].

The gestational period of the participants showed that those in their 2nd trimester have the highest prevalence with 25(56.8%). This is consistent with the result reported by Okonko, [16]. This difference may be as a result of either change in urinary stasis and vesicoureteral reflux or decrease in urinary progesterones and oestrogens in the various trimester of pregnancy.

Based on level of education, the highest prevalence was among those who attained primary level of education with 8(88.9%), while those with tertiary education had 28.9%, difference could be attributed to awareness and hygiene. Also based on marital status of participants, the unmarried status had a distributions of 8(72.7%), while the married status had 35 (25.2%) with UTI. This could be due to the level of socioeconomic tendencies of the participants.

Table 1. Prevalence of urinary tract infections in pregnant women in relation to age

Age group	Frequency	Number positive	(%)	P-value
20-25	41	7	17.1	0.006
26-31	69	26	37.7	
32-37	25	8	32	
38-43	13	2	15.4	
44-47	2	0	0	
Total	150	43		

Table 2. Distribution of urinary tract infections in pregnant women in relation to gestational period

Gestation (week)	Frequency	Number positive	(%)	p- value
1-12	22	3	13.6	< 0.0001
13-25	84	15	17.9	
26-40	44	25	56.8	
Total	150	43	100	

Table 3. Distribution of urinary tract infections in relation to other factors

Factors		N	NOS POS	(%)	p value
Status	Married	139	35	25.2	0.126
	Unmarried	11	8	72.7	
Education	Primary	9	8	88.9	0.0001
	Secondary	72	15	20.8	
	Tertiary	69	20	28.9	
Past history of UTI	Present	101	33	32.7	129
	Absent	49	10	20.4	
Sexual activity	Active	135	42	31.1	0.067
	Non-active	15	1	6.7	
Haemoglobin genotype	AA	114	38	33.3	0.683
	AS	33	5	15.2	
	AC	3	0	0	

Table 4. Distribution OF UTI in prenant women in relation to parity

Parity	NOS examined	NOS with UTI	(%)	P-value
0	32	10	31.3	0.179
1	57	12	21.1	
2	39	6	15.4	
3	36	8	22.2	
>4	16	7	43.8	
Total	150	43		

Table 5. Distribution of isolates from pregnant women

Isolates	Number isolated	(%)
Bacillus specie	11	8.5
S. epidermidis	11	8.5
Escherichia coli	7	5.4
Enterobacter specie	7	5.4
S. aureus	46	35.7
Lactobacillus specie	3	2.3
Candida albicans	35	27.1
Klebsiella aerogenes	9	6.9
Total	129	

Among isolated pathogens, *Staphylococcus aureus* was the most predominant bacteria with 46(35.7%).”This is in contrast to other results gotten in other studies conducted in Babol city, *E. coli* was indicated to be the cause of 83% of UTIs in pregnant women [17] and Moreover, in studies conducted by Masinde in Tanzania, Al-Haddad in Yemen, Hamdan in Sudan and Totsika in Australlia [18], *E. coli* was the main cause of UTIs among women of those areas and the prevalence in those regions were 47.2%, 41.5%, 42.4%, respectively”.

5. CONCLUSION

This study concludes that relationship between UTI and parity is not statistically significant. However, education status and gestational growth are statistically significant in relation to UTI among pregnant women, regardless of other associated risk factors such as haemoglobin genotype, past history of UTI and marital status.

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

This study was approved by the ethical committee of Igbinedion University Teaching Hospital, Okada Edo state. The purpose of this study was well explained to the pregnant women and strict adherence to confidentiality maintained.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Ezugwu IE, Afunwa RE, Onyia FA, Chukwunwejim CR, Eze AE. Prevalence of urinary tract infections and associated risk factors among pregnant women in Enugu Metropolis. *Nigerian Journal of Biosciences and Medicines*. 2021;9(10).
2. Ali M, Abdallah MS. Prevalence of urinary tract infection among pregnant women in Kano, Northern Nigeria. *Archives of Reproductive Medicine and Sexual Health*. 2019;2(1):23-29.
3. Tamalli M, Sangar B, Alghazal AM. Urinary tract infection in pregnancy at Al-khoms, Libya. *International Journal of Medicine and Medical Sciences*. 2013;3(5):455-459.
4. Nwachukwu E, Onyebuchi O, Michael O. Prevalence of urinary tract infections in pregnant women in Onitsha, Nigeria. *Journal of Bacteriology Mycology*. 2018;6(5):284–285. DOI: 10.15406/jbmoa.2018.06.00219
5. Giraldo PC, Araújo ED, Junior JE, Amaral GD, Passos RL, Gonçalves AK. The prevalence of urogenital infections in pregnant women experiencing preterm and full-term labor. *Infectious Diseases Obstetrics Gynecology*. 2012:1–4. DOI: 10.1155/2012/878241
6. Taye, S., Getachew, M., Desalegn, Z., Biratu, A., & Mubashir, K. Bacterial profile, antibiotic susceptibility pattern and associated factors among pregnant women with urinary tract infection in Goba and Sinana Woredas, bale zone, Southeast Ethiopia. *BMC Research Notes*. 2018;11(1):799.
7. Nabbugodi WF, Gichuhi JW, Mugo NW. Prevalence of urinary tract Infection, microbial etiology, and antibiotic sensitivity pattern among antenatal women presenting with lower abdominal pains at Kenyatta National Hospital, Nairobi, Kenya. *The Open Access Journal of Science and Technology*. 2015;3:1-6.
8. Cheesbrough M. *District Laboratory practice in tropical countries (Part II)*. Cambridge University. 2010:19-11.
9. Fawole MO, Oso BA. *Laboratory manual for microbiology*. Spectrum Book Ltd, Ibadan. 5th edition. 1988;1:22-45.
10. Holt JG, Krieg NR, Sneath PHA, Stanley JT, Williams ST. *Bergey's manual of determinative bacteriology*, 3rd edition. Williams and Wilkins, Baltimore. 1994;9:787.
11. Cowan ST, Steel KJ. *Cowan and steel's manual for the identification of medical bacteria*. (5th Edn). Cambridge University Press, Cambridge, London. 2003:1-210.
12. El-Kashif MML. Urinary tract infection among pregnant women and its associated risk factors: A cross-sectional study. *Biomedical Pharmacology Journal*. 2019;12(4):2003–2010.
13. Thakur S, Nagpal KL. Urinary tract infection in pregnant women at Kathmandu, Nepal. *Journal of Medical Care Research Review*. 2020;3(9):454–458.

14. Mobbasheri ET, Ghaemi E, Moujloo M, Vakili MA. Prevalence of bacteriuria during pregnancy in Gorgan, Iran. Gorgan Medical Journal. 2001;9:42–47.
15. Gibbs RS, McGregor JA, Mead PB, Eschenbach DA, Hager WD, Sweet RL. A survey of practices in infectious diseases by obstetrician-gynecologists. Obstetrics and Gynecology. 1994;83(4):631–636.
16. Okonko IO, Ijandipe LA, Ilusanya OA, Donbraye-Emmanuel OB, Ejembi J, Udeze AO, Egun OC, Fowotade A, Nkang AO. Incidence of Urinary Tract Infection (UTI) among pregnant women in Ibadan, South-Western Nigeria. African Journal of Biotechnology. 2009;8(23): 6649-6657.
17. Yaghibi RH, et al. The role of hygienic behavior in incidence of urinary tract infection among pregnant women referred to health center, Babol, 2003-2005. Journal of Babol University Medical Science. 2006;8(2):56–62.
18. Totsika M, Moriel DG, Idris A, Rogers BA, Wurpel DJ, Phan MD. Uropathogenic *Escherichia coli* mediated urinary tract infection. Current Drug Targets. 2012; 13(11):1386–1399.

© 2023 Sule et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95686>