

Unusual Effects of Intramolecular Hydrogen Bonds and Dipole Interaction on FTIR and NMR of Some Imines

Elham Abdalrahem Bin Selim¹ and Mohammed Hadi Al-Douh^{1*}

¹*Department of Chemistry, Faculty of Science, Hadhramout University, Mukalla, Hadhramout, Republic of Yemen.*

Authors' contributions

This work was carried out in collaboration between both authors. Author MHAD designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author EABS managed the analyses of the study. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJACR/2019/v3i330096

Editor(s):

(1) Dr. Endang Tri Wahyuni, Professor, Department of Chemistry, Gadhah Mada University, Indonesia.

Reviewers:

(1) A. Ayeshamariam, Khadir Mohideen College, India.

(2) A. Veera Reddy, Suven Life Sciences, India.

(3) Pipat Chooto, Prince of Songkla University, Thailand.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50408>

Case Study

Received 15 May 2019
Accepted 23 July 2019
Published 16 August 2019

ABSTRACT

Unusual effects of intramolecular hydrogen bonds and dipole interactions are investigated using FTIR, NMR and X-Ray crystallography analyses of some imines. These phenomena affect both FTIR absorptions and chemical shifts.

Keywords: Intramolecular hydrogen bonds; dipole interactions; FTIR; ¹H NMR.

1. INTRODUCTION

Organosynthetic chemistry field is important in the drug discovery, which needs some of the different analytical characterizations for identification and confirmation of the chemical

structure of these compounds. A hydrogen atom in organic compounds that is bonded to oxygen O or nitrogen N can participate in hydrogen bonding. This electrophilic hydrogen can form intramolecular and intermolecular attachments with nonbonding electrons on O or N atoms.

*Corresponding author: Email: mhd_douh@yahoo.com, mhdouh@hu.edu.ye;

A hydrogen bond is a strong dipole-dipole attraction. In some organic compounds, we can see that intramolecular hydrogen bonding and dipole interactions have a strong effect on the elongation of N–H and O–H bonds resulting in unusual values of FTIR and ^1H NMR of some organic compounds, as shown in compounds 1–5.

The X-Ray crystallographic analyses of 1 and 2 were determined, and the intramolecular hydrogen bonds of both imines were compared. The length of the intramolecular hydrogen bonds, $-\text{C}=\text{N}\cdots\text{H}-\text{O}$ and $-\text{C}=\text{N}\cdots\text{H}-\text{N}$ in 1 are at the respective 1.77 and 2.23 Å [1]. So, the bond $-\text{C}=\text{N}\cdots\text{H}-\text{N}$ form weaker intramolecular hydrogen bonds than the bond $-\text{C}=\text{N}\cdots\text{H}-\text{O}$ because the nitrogen atom is less electronegativity than oxygen atom. While the intramolecular hydrogen bond $-\text{C}=\text{N}\cdots\text{H}-\text{N}$ in 2 has two values (2.31 and 2.29 Å) for both stereoisomers of 2 [2]. However, imines 1 and 2 are used in synthesise of bis-Schiff base [3] and benzimidazole derivatives [4–6], which are used widely in pharmaceutical and biological applications [7–9].

2. CHEMICAL STRUCTURE AND ITS APPLICATION

The effect of N–H and O–H intramolecular hydrogen bonding can be seen in the case of an imine 1 as shown in Fig. 1. The FTIR spectrum of this compound exhibits only one spike around 3414 cm^{-1} for a single N–H stretching bond instead of two spikes of a primary amino group as in Fig. 2. This may be assigned to the intramolecular hydrogen bonding of nitrogen atom in $\text{N}=\text{C}-\text{H}$ and one hydrogen atom in the

amino group. On another hand, the chemical shifts of hydroxyl proton and a proton that attached to carbon number seven both appear as a singlet at $\delta = 13.56$ and 8.63 ppm, respectively, Fig. 3 [10].

These unexpected values can be explained by intramolecular hydrogen bonding which generates five and six ring motifs as in Fig. 1. Greater deshielding leads to unusual delta values as in Fig. 3. Imine 2 forms stronger dipole-dipole interaction. As a result, the chemical shift of a proton that attached to carbon number seven appears at $\delta = 8.79$ ppm, which is uncommon as shown in Fig. 4, similar to that from previous studies [11].

The effect of hydrogen bonds reflects on the FTIR absorptions of diazo dyes 3, 4 and 5, Fig. 5. It is believed the lower symmetrical stretching of nitro groups at *ortho* position in 3 which appears at 1301 cm^{-1} due to the intramolecular hydrogen bonding and the dipole interaction between them with phenolic and aldehydic hydrogen's atoms, Fig. 5 [12].

These overlaps leave these bonds to have a little electron density and a little less stiff. As a result, they vibrate a little more slowly. The O–H stretching frequency in 3 reflects the stronger intramolecular hydrogen bonding as a very broad absorption around 3277 cm^{-1} shown in Fig. 6. However, the IR spectra of both 4 and 5 display slightly lower asymmetrical stretching frequencies of $-\text{NO}_2$ and O–H due to intramolecular and intermolecular hydrogen bonding, respectively as in Fig. 6 and the results previously reported [12].




Fig. 1. Imines 1 and 2




Fig. 2. FTIR spectra of imines 1 and 2




Fig. 3. ¹H NMR spectrum of 1




Fig. 4. ^1H NMR spectrum of 2




Fig. 5. The dipole interaction and intramolecular hydrogen bonding in both 3 and 4, and intermolecular hydrogen bonding in 5




Fig. 6. The FTIR spectra of diazo dyes 3, 4 and 5

3. CONCLUSION

In this work, we have reported some imines that have strong effects on intramolecular hydrogen bonding and dipole interaction which leads to the unexpected FTIR absorptions and the unusual values of chemical shifts in ¹HNMR.

ACKNOWLEDGEMENTS

We thank Mrs. Gehan A. Balala in PCB Research Laboratory, Faculty of Science, Hadhramout University for FTIR analyses.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Al-Douh MH, Hamid SA, Osman H, Kia R, Fun HK. 2-Amino-N-(2-hydroxy-3-methoxybenzylidene) aniline. *Acta Crystallographica Section E*. 2008;64(7): 1201–1202. DOI: 10.1107/S1600536808016292.
2. Al-Douh MH, Hamid SA, Osman H, Kia R, Fun HK. 2-Amino-N-(2-benzyloxy-3-methoxybenzylidene) aniline. *Acta Crystallographica Section E*. 2008;64(7): 1290–1291. DOI: 10.1107/S1600536808017844.
3. Al-Douh MH, Hamid SA, Osman H. Synthesis, spectroscopic and crystal structures studies of N,N'-bis(2-hydroxy-3-methoxybenzylidene) diaminobenzene derivatives. *University of Aden Journal of Natural and Applied Sciences*. 2010;14(2): 325–347. [ISSN: 1606-8947]
4. Al-Douh MH, Osman H, Hamid SA. Synthesis and characterization of N-1-(2-hydroxy-3-methoxybenzyl)-2-(2-hydroxy-3-methoxyphenyl)-1H-benzimidazole. *Hadhramout University Journal of Natural and Applied Sciences*. 2011;8(1):11–32. [ISSN: 2227-9164]
5. Al-Douh MH, Osman H, Hamid SA. Synthesis and characterization of novel benzimidazole: N-1-(2-benzyloxy-3-methoxybenzyl)-2-(2-benzyloxy-3-methoxyphenyl)-1H-benzimidazole. *University of Aden Journal of Natural and Applied Sciences*. 2011;15(2):375–393. [ISSN: 1606-8947]
6. Al-Douh MH, Osman H, Hamid SA. Synthesis and characterization of 2-(2-hydroxy-3-methoxyphenyl)-1H-benzimidazole. *Hadhramout University Journal of Natural and Applied Sciences*. 2012;9(1):49–64. [ISSN: 2227-9164]
7. Al-Douh MH, Sahib HB, Osman H, Hamid SA, Salhimi SM. Anti-proliferation effects of benzimidazole derivatives on HCT-116 colon cancer and MCF-7 breast cancer cell

- lines. Asian Pacific Journal of Cancer Prevention. 2012;13(8):4075–4079.
DOI: 10.7314/APJCP.2012.13.8.4075.
8. Helal MH, Al-Douh MH, Hamid SA, Osman H, Salhimi SM, Majid AM. Diaminobenzene Schiff base induces caspase 9-dependent apoptosis in U937 Leukemia cells. Webmedcentral. 2010; WMC001081:1–13.
[ISSN: 2046-1690]
Available: http://www.webmedcentral.com/article_view/1081.
 9. Helal MH, Al-Mudaris ZA, Al-Douh MH, Osman H, Wahab HA, Al-Najjar BO, Abdallah HH, Majid AM. Diaminobenzene Schiff base, A novel class of DNA minor groove binder. International Journal of Oncology. 2012;41(2):504–510.
DOI: 10.3892/ijo.2012.1491.
 10. Al-Douh MH, Osman H, Hamid SA. Synthesis, NMR spectroscopy and crystal structures studies of 2-amino-N-(2-hydroxy-3-methoxybenzylidene) aniline. Proceeding of International Conference on Instrumentation, Communication, Information Technology and Biomedical Engineering (ICICI BME 2009). Aula Timur, Bandung Institute of Technology, Ganesha 10, Bandung 40132, Indonesia. 2009;247-253.
[ISBN 978-979-1344-67-8]
IEEE CFP0987H-CDR
 11. Al-Douh MH, Osman H, Hamid SA. Synthesis, NMR spectroscopy and crystal structures studies of novel 2-amino-N-(2-benzyloxy-3-methoxybenzylidene) aniline. Proceeding of International Conference on Instrumentation, Communication, Information Technology and Biomedical Engineering (ICICI BME 2009). Aula Timur, Bandung Institute of Technology, Ganesha 10, Bandung 40132, Indonesia. 2009;438-443.
[ISBN 978-979-1344-67-8]
IEEE CFP0987H-CDR
 12. Al-Douh MH, Bin-Selim EA, Al-Haik WM, Bin-Mahram SM, Al-Bakri AK, Al-Nohey DS, Sonbol KS. Synthesis and antimicrobial activity of some compounds derived from diazo dyes. International Journal of Chemistry Studies. 2019;3(4):6-12.
[ISSN: 2581-348X]

© 2019 Selim and Al-Douh; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50408>