



Role of Genetic Variability and Their Associated Character on Yield Attributing Traits in Castor (*Ricinus communis* L.) Germplasm Lines

Yenametla Rajavardhan ^a, J. Jawahar Lal ^{b*},
P. Lakshamma ^b, C. V. Sameer Kumar ^a and A. Anil Kumar ^b

^a Department of Genetics and Plant Breeding, College of Agriculture, PJTSAU, Hyderabad (Telangana), India.

^b Indian Institute of Oilseeds Research, Hyderabad, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i102635

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104098>

Original Research Article

Received: 28/05/2023

Accepted: 03/08/2023

Published: 14/08/2023

ABSTRACT

Castor (*Ricinus communis* L.) is an industrially important non-edible oil seed crop with limited genetic variation. It is consequently important to diversify the genetic base to utilize the diversity. The primary objective of this study was to assess the extent and significance of variability among castor genotypes concerning yield and its associated traits. A study with two hundred germplasm lines along with four checks on genotypes on variability, correlation, and path coefficient was carried out for nine traits. The genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) values indicated the presence of broad variation for all characters except for days to 50 percent flowering, days to maturity, number of nodes, and oil content. However, high heritability coupled with high genetic advance as percent mean was observed for plant height, days

*Corresponding author: E-mail: jawaharlaljatoth@gmail.com;

to 50 percent flowering, days to maturity, number of nodes, total length of the primary spike, number of effective spikes, seed weight, and total yield. The character's plant height, number of nodes, total length of the primary spike, number of effective spikes, seed weight, and oil content show a positive significant correlation with the total yield. The path coefficient analysis indicates that the utmost positive direct effect on total yield was exerted by a number of effective spikes per plant. Hence it was concluded that selection for traits total length of the primary spike, number of effective spikes, and 100 seed weight beneficial in yield improvement programs of castor.

Keywords: Castor; variability; correlation; path coefficient; yield.

1. INTRODUCTION

“Castor (*Ricinus communis* L.) is an industrially important non-edible oil crop that belongs to the family of Euphorbiaceae and the genus *Ricinus*. It is a diploid plant with a chromosome number of $2n=20$ ” [1]. Worldwide castor is cultivated in an area of 1296.9 ('000 ha) with a production of 1396 ('000 t). Globally, around 30 countries are cultivating castor on a commercial scale and among them India, China, Brazil and USSR are the major castor producer [2,3]. India is leading in the production and export of castor in the world [4]. “Castor seeds contain 45 to 55 percent oil which is the major source of ricinoleic acid an unusual hydroxyl fatty acid that has tremendous industrial significance” [5]. The effectiveness of crop improvement initiatives depends on the type and extent of genetic variability present in the crop. To accomplish this, the genotypes must undergo phenotypic assessment, followed by the estimation of trait heritability and their interrelationships. Heritability estimates are vital for discerning the heritable component of variability within the traits. These steps are essential for gaining insights into the heredity aspects and determining the potential for improvement in the crop [6]. To make selection effective the heritability should couple with high genetic advance [1]. To achieve the desired yield improvement, it is imperative to genetically enhance traits contributing to yield. Correlation coefficient studies offer valuable insights into the nature and strength of associations among different traits, while path coefficient analysis helps identify direct and indirect effects on other attributes through correlations, establishing causal relationships between variables. These methods play a crucial role in understanding the interplay of traits and guiding targeted crop enhancement strategies [7]. Hence the present investigation was undertaken to estimate the of genetic variability and character associations among yield-related traits in the selected castor germplasm lines.

2. MATERIALS AND METHODS

The experiment was conducted at the Indian Institute of Oilseeds Research (IIOR), Rajendranagar, Hyderabad. The city is nestled on the Deccan Plateau and is positioned at a $17^{\circ}15'16''$ N latitude and $78^{\circ}18'30''$ E longitude during *Kharif*, 2022. Two hundred germplasm lines along with four checks (DCS-9, GC-3, 48-1, PCS-262) with a spacing of 90 cm between the rows and 60 cm between the plants were evaluated in an augmented randomized complete block design by replicating the checks in which each line is sown in a single row of 6.0 m length and all recommended agronomic practices adopted to raise a good crop.

The observations viz., Plant height (cm), days to 50% flowering, days to maturity of primary spike, number of nodes to primary spike, total length of primary spike(cm), number of effective spikes per plant, 100 Seed weight(g), total seed yield/plot(g), oil content (%) were recorded on randomly selected five plants in each genotype and mean values were considered for analysis.

An augmented randomized complete block design [8] was employed to assess genotypic differences among the entries, following the method suggested by Panse and Sukhatme [9]. Genotypic (σ^2_g) and phenotypic (σ^2_p) variances were calculated using Burton and De vane's formula [10], and the variation range was categorized based on Subramanian and Menon's recommendations [11]. Broad sense heritability [h^2 (b)] was determined using the formula by Hanson [12], and heritability estimations were classified according to Johnson et al. [13]. Genetic advance, indicating the projected gain in the next generation, was calculated using the formula of Johnson et al, [13]. Variance and covariance components for each pair of characteristics were estimated using the methods of Al-Jibouri et al, [14] to calculate

phenotypic and genotypic correlation coefficients. The significance of correlation coefficients was tested by comparing the estimated values with table values [15] at 5 percent and 1 percent levels. Path coefficient analysis, as proposed by Wright [16] and detailed by Dewey and Lu [17], was used to distinguish the direct and indirect effects of independent factors on the dependent variable. Analysis was performed by using the INDOSTAT method.

3. RESULTS AND DISCUSSION

3.1 Genetic Variability Parameters

Analysis of variance revealed the significant variation among the genotypes for all the traits studied in two hundred germplasm lines. The values of *gcv* and *pcv* were categorized as low (20%) [11]. The *gcv* values ranged from 7.03 for oil content to 48.02 for the number of effective spikes (Table 1). High *gcv* values were recorded for the traits plant height (28.66), total length of the primary spike (25.38), number of effective spikes (48.02), 100 seed weight (28.94), total yield (34.20) and medium values for the characters days to 50 percent flowering (19.46), days to maturity (13.46), number of nodes (15.53) and oil content (7.03) shows the low *gcv* value. High *pcv* values for the characters plant height (28.78), total length of primary spike (25.56), number of effective spikes (48.68), seed weight (29.36), total yield (34.20), and medium values for the traits days to 50 percent flowering (19.72), days to maturity (13.66), number of nodes (16.88) and oil content (7.32) shows the

low *pcv* value. *gcv* values were slightly lower than those of *pcv* indicating the minor influence of the environment on the expression of these characters. Results also show the presence of broad genetic variability for all the traits under study except oil content. Similar results can be observed in the findings of Movaliya et al. [18]; Chaudhari et al. [19]; Mullualem et al. [1]; Alhaji et al. [20].

High heritability was found for all the characters under the study with the maximum heritability found in the trait total yield (Table 1). However, only eight among the nine characters studied viz., plant height, days to 50 percent flowering, days to maturity, number of nodes up to primary spike, the total length of the primary spike, number of effective spikes, seed weight, and total yield coupled shows the high heritability coupled with the high genetic advance as percent of mean which indicated a preponderance of additive genetic effect for the characters and there was no environmental influence on the expression and selection for improvement of such characters could be rewarding. Oil content shows high heritability coupled with moderate genetic advance as a percent mean which indicated the presence of both additive and non-additive gene effects and considerable effect of the environment. "High heritability accompanied by moderate genetic advance as a percent mean indicated that the genotypes, under study, were diverse with immense genetic potential and a further improvement in these traits is possible by practicing a simple selection technique" [18,21].

Table 1. Magnitude of variability, heritability, and genetic advance as percent of mean for different characters in the germplasm lines of castor studied during *kharif*, 2022

Trait	Mean	Range		GCV	PCV	h_{BS}	GAM
		Minimum	Maximum				
PH	94.96	36.80	178.00	28.66	28.78	99.16	58.87
DFF	68.97	43.00	104.00	19.46	19.72	97.45	39.64
DM	117.19	79.00	160.00	13.46	13.66	97.19	27.38
NN	17.27	10.00	24.60	15.53	16.88	84.63	29.47
TLPS	28.43	14.00	52.00	25.38	25.56	98.58	51.99
NES	12.94	4.00	28.00	48.02	48.68	97.32	97.74
SW	33.10	8.30	59.80	28.94	29.36	97.20	58.87
OC	44.80	27.55	51.46	7.03	7.32	92.42	13.95
TY	815.48	267.4	1698.9	34.01	34.20	98.91	68.78

PH - Plant height up to primary raceme (cm), DFF - Days to 50 percent flowering, DM - Days to maturity, NN - Number of nodes up to primary raceme, TLPS - Total length of the Primary spike (cm), NES - Number of effective spikes per plant, SW - 100 seed weight (g), OC - Oil content (%) and TY - Total Seed yield per plot (g). PCV - Phenotypic coefficient of variation, GCV - Genotypic coefficient of variation, h_{BS} - Heritability at broad sense, GAM - Genetic advance as a percent mean

3.2 Character Association Studies

Seed yield is a multifaceted trait influenced by numerous characteristics. Breeders seek genetic diversity in these traits to identify favourable types. In this investigation, correlations and path coefficient analyses were employed to explore the associations among yield-related traits and their connection to overall yield.

Yield is a multifaceted trait influenced by various contributing characters that can exhibit positive or negative associations both with yield and among themselves. The extensive correlations computed using the Pearson correlation coefficient offer valuable insights for indirect selection strategies to enhance overall yield. The results of the Pearson correlation analysis depict relationships between yield and its contributing characters (Table 2, Fig. 1).

The results revealed that the traits plant height (0.2505^{**}), number of nodes up to primary spike (0.4202^{**}), total length of the primary spike (0.3451^{**}), number of effective spikes (0.8470^{**}), seed weight (0.6301^{**}), oil content (0.4964^{**}) have the significant positive correlation with the seed yield indicating that these attributes were influencing more on seed yield in castor and therefore important for bringing improvement in seed yield. A non-significant positive correlation was registered for the days to maturity (0.0394) while days to 50 percent exhibited a non-significant negative correlation with seed yield which is highly useful in developing short-

duration varieties that require negative selection for flowering without compromising the seed yield. These kinds of correlations were similar to the findings of Movaliya et al. [18]; Ruksar et al. [21]; Reddy et al. [22].

To gain a comprehensive understanding and enable effective manipulation of the characters, path coefficients were employed alongside correlation coefficients. Path analysis allows for the dissection of correlations into direct and indirect effects, providing insights into the relative significance of the causal factors involved [17]. In the present study, nine characters were taken and partitioned into direct and indirect effects using seed yield per plot as a dependent variable. The information about the direct and indirect effects of eight variables on seed yield per plot in two hundred germplasm lines castor is described below (Table 3, Fig. 2).

The number of effective spikes per plant (0.7136), seed weight (0.1356), and total length of the primary spike have a high positive direct effect on seed yield while oil content (0.0542), days to maturity (0.0478) and plant height (0.0341) have a low positive direct effect on seed yield. Due to the positive direct effect on seed yield these traits can be considered for the crop improvement program of castor. Days to 50 percent flowering (-0.0622) and number of nodes (-0.0117) have negative direct effects due to negative indirect effects of other traits. Patel et al. [24]; Rukhsar et al. [21]; Alhaji et al. [20] findings were similar to the present study.




Fig. 1. Phenotypic correlation coefficients among different traits in castor

Table 2. Correlation coefficients of yield attributing traits in the germplasm lines of castor studied during *kharif*, 2022

	Plant Height (cm)	Days to 50 % Flowering	Days to Maturity	Number of Nodes to primary raceme	Total Length of the Primary Spike (cm)	Number of Effective Racemes	100 seed weight (g)	Oil content (%)	Seed Yield (g)
Plant Height (cm)	1	0.2924**	0.2739**	0.4646**	0.2373**	0.1773*	0.3954**	0.3187**	0.2505**
Days to 50 % Flowering		1	0.6395**	0.3970**	-0.0624	-0.0363	0.2002**	-0.1189	-0.0394
Days to Maturity			1	0.2873**	0.0008	0.0015	0.1745*	-0.0784	0.0345
Number of Nodes to primary raceme				1	0.1922**	0.4424**	0.5424**	0.2548**	0.4202**
Total Length of the Primary Spike (cm)					1	0.2086**	0.3346**	0.3051**	0.3451**
Number of Effective Racemes						1	0.5893**	0.4475**	0.8470**
100 seed weight (g)							1	0.5402**	0.6301**
Oil content (%)								1	0.4964**
Seed Yield (g)									1

** Significant at 0.01; *Significant at 0.05

Table 3. Path coefficient analysis of agro morphological traits in the germplasm lines of castor studied during *kharif*, 2022

	Plant Height (cm)	Days to 50 % Flowering	Days to Maturity	Number of Nodes to primary raceme	Total Length of the Primary Spike (cm)	Number of Effective Racemes	100 seed weight (g)	Oil content (%)
Plant Height (cm)	0.0341	0.0100	0.0093	0.0158	0.0081	0.0060	0.0135	0.0109
Days to 50 % Flowering	-0.0182	-0.0622	-0.0398	-0.0247	0.0039	0.0023	-0.0125	0.0074
Days to Maturity	0.0131	0.0305	0.0478	0.0137	0.0000	0.0001	0.0083	-0.0037
Number of Nodes to primary raceme	-0.0054	-0.0047	-0.0034	-0.0117	-0.0023	-0.0052	-0.0064	-0.0030
Total Length of the Primary Spike (cm)	0.0296	-0.0078	0.0001	0.0240	0.1246	0.0260	0.0417	0.0380
Number of Effective Racemes	0.1265	-0.0260	0.0011	0.3157	0.1489	0.7136	0.4206	0.3194
100 seed weight (g)	0.0536	0.0272	0.0237	0.0736	0.0454	0.0799	0.1356	0.0733
Oil content (%)	0.0173	-0.0064	-0.0043	0.0138	0.0165	0.0243	0.0293	0.0542
Seed Yield (g)	0.2505	-0.0394	0.0345	0.4202	0.3452	0.8470	0.6302	0.4964
Partial R ²	0.0085	0.0025	0.0016	-0.0049	0.0430	0.6044	0.0855	0.0269

R Square = 0.7675 Residual effect = 0.4822




Fig. 2. Phenotypal path diagram of various traits with seed yield per plot in castor

4. CONCLUSION

The current investigation on variability parameters, correlation, and path coefficient analyses revealed that the characters length of the primary spike, number of effective spikes, and 100 seed weight largely define the seed yield of the castor. Therefore, it is beneficial to put more focus on these traits in the selection process in order to develop high-yielding varieties and hybrids castor.

5. FUTURE SCOPE

Through extensive research on GCV (genotypic coefficient of variation), PCV (phenotypic coefficient of variation), heritability, genetic advance as a percentage of mean, correlation, and path coefficient analysis, successfully identified the most desirable traits for enhancing yield in castor. These findings offer valuable insights that can be leveraged to enhance seed yield without compromising other important characteristics such as crop duration and plant architecture in future castor breeding programs. Moreover, the gathered data holds significant potential for selecting superior lines in the development of improved varieties and hybrids, fostering advancements in castor cultivation.

ACKNOWLEDGEMENT

This is part of the author's post-graduate thesis work at Professor Jayashankar Telangana State

Agricultural University. The author is highly grateful for the research facilities provided by the Indian Institute of Oilseeds Research, Hyderabad, and Professor Jayashankar Telangana State Agricultural University.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mullualem D, Alemaw G, Petros Y, Alemu S. Phenotypic variability and association of traits among yield and yield-related traits in Castor (*Ricinus communis* L.) accessions at Melkassa, Central Rift Valley of Ethiopia. African Journal of Agricultural Research. 2017;12(52):3562-3568.
2. Damodaram T, Hegde DM. Oilseeds situation: A statistical compendium. Directorate of Oilseeds Research, Hyderabad. 2010;486.
3. Rajput D, Patel MP, Kumar S, Patel R, Katba P. Screening of castor germplasm for wilt reaction and morpho-molecular characterization of resistant genotypes. Heliyon. 2023;9(3).
4. Sadaiah K, Neelima G, et al. Genetic parameters, diversity and character association studies in germplasm lines of castor (*Ricinus communis* L.). Electronic

- Journal of Plant Breeding. 2021;12(4):1134-1141.
5. Senthilvel S, Shaik M, Anjani K, Shaw RK, Kumari P, Sarada C, Ushakiran B. Genetic variability and population structure in a collection of inbred lines derived from a core germplasm of castor. *Journal of Plant Biochemistry and Biotechnology*. 2016;26(1):27-34.
 6. Jyothsna S, Patro TSSK, Rani YS, Neeraja B, Ashok S, Triveni U. Studies on genetic parameters, character association, and path analysis of yield and its components in Little Millet (*Panicum sumatrense*). *International Journal of Agriculture Sciences*, 2016;8(5):1018-1020.
 7. Dapke JS, Naik MR, Singh AP, Baraskar VV, Narwade AV, Indivar P. Genetic association yield with different agronomic traits in castor (*Ricinus communis* L.). *International Journal of Biology Research*. 2016;1(1):47-49.
 8. Federer WT. Augmented (or Hoonuiaku) designs. *Hawaiian planers record*. 1956;55:191-208.
 9. Panse VG, Sukhatme PV. *Statistical methods for agricultural workers*, ICAR, New Delhi; 1985.
 10. Burton CW, De Vane EH. Estimating heritability in tall festuca (*restuhaarundinacae*) from replicated clonal material. *Agronomy Journal*. 1953;45:1476-1481.
 11. Subramanian PS, Menon PM. Genotypic and phenotypic variability in Rice. *Madras Agricultural Journal*. 1973;60:1093.
 12. Hanson WD. *Heritability, statistical genetics, and plant breeding* NASNRC. Washington Publishers. 1956;98:115-140.
 13. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in Soybean. *Agronomy Journal*. 1955;47:314-318.
 14. Al-Jibouri HA, Miller PA, Robinson HF. Genotypic and environmental variation and correlation in upland cotton cross of interspecies origin. *Agronomy Journal*. 1958;50:633-636.
 15. Fisher RA, Yates F. *Statistical tables for biological, agricultural and medical research*, Hafner, New York. 1963;6:55-56.
 16. Wright S. *Correlation and causation*. *Journal of Agricultural Research*. 1921;20:557-585.
 17. Dewey JR, Lu KH. Correlation and path coefficient analysis of components of crested wheat grass seed production. *Agronomy Journal*. 1959;51:515-518.
 18. Movaliya HM, Chovatia VP, Madariya RB, Mungala RA, Pipaliya HR, Bhuva SK. Study of variability and correlation for seed yield and its attributes in castor (*Ricinus communis* L.). *Journal of Pharmacognosy and Photochemistry*. 2018;7(2):1474-1477.
 19. Chaudhari D, Parmar V, Dube D, Bhakta R, Patel A, Lodam V. Genetic variability and divergence study in castor (*Ricinus communis* L.). *Advances in Life Science*. 2016;5(16):6418-6422.
 20. Alhaji KA, Oliseh A, Salihu BZ, Kabaraini MA. Studies on genetic variability and trait relationship in castor (*Ricinus communis* L.). *Journal of Plant Development*. 2019;26:3-11.
 21. Rukhsar Patel MP, Parmar DJ, Kumar S. Genetic variability, character association, and genetic divergence studies in castor (*Ricinus communis* L.). *Annals of Agrarian Science*. 2018;16(2):143-148.
 22. Reddy Sahithi P, Bindu Priya P, Sadaiah K, Vanisri S, Sameer Kumar CV. Genetic variability and character association studies among yield attributing traits in genepool lines of castor (*Ricinus communis* L.). *Biological Forum – An International Journal*. 2022;14(2):1277-1281.
 23. Patel JK, Nakarani DB. Character association and path analysis in castor (*Ricinus communis* L.). *International Journal of Agricultural Sciences*. 2016;12(1):22-27.

© 2023 Rajavardhan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104098>