



# **Effect of Foliar Application of Micronutrients on Growth, Yield and Flower Quality of Gerbera (*Gerbera jamesonii*) under Naturally Ventilated Polyhouse Conditions in Prayagraj**

**Sunkara Mounika<sup>a++\*</sup> and Urfi Fatmi<sup>a#</sup>**

<sup>a</sup> *Department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh, India.*

## **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/IJECC/2023/v13i102670

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104662>

**Original Research Article**

**Received: 07/06/2023**

**Accepted: 11/08/2023**

**Published: 17/08/2023**

## **ABSTRACT**

A field experiment was executed during November, 2022 to April, 2023 under polyhouse conditions in the Research Field, Department of Horticulture, SHUATS, Prayagraj. The objective was to study the response of foliar application of micronutrients on growth, yield and flower quality of gerbera. The research consisted of three levels of zinc, iron, magnesium (0.2, 0.4, 0.6%). The experiment was laid out in Randomized Block Design with ten treatments and replicated thrice. The results indicated that the treatment T2-Zn@0.4% performed better in all parameters recorded viz., plant height (24.08 cm), plant spread (37.68 cm), days for first bud initiation (73 days), number of flowers per plant (10), stalk length (54.17 cm) and vase life (9 days).

<sup>++</sup>M.Sc. Scholar;

<sup>#</sup>Assistant Professor;

\*Corresponding author: E-mail: [mounikasunkara55@gmail.com](mailto:mounikasunkara55@gmail.com);

**Keywords:** Micronutrient; iron; magnesium; zinc; gerbera; growth; yield.

## 1. INTRODUCTION

Gerbera (*Gerbera jamesonii* Bolus) is an important commercial cut flower grown throughout the world in a wide range of climatic conditions. It belongs to the family Asteraceae. It is commonly known as Transvaal Daisy, Barberton Daisy or African Daisy and this flower is native to South African and Asiatic regions. The world consumption of flowers is estimated to be US dollar 60 billion per annum in which India contribute 0.38 percent only [1]. The estimated area under flower cultivation in India is about 0.90 thousand ha with a production of 21.84 cut flowers in lakh number and 4.00 thousand metric tonnes as a loose flower. It ranks fourth in the international cut flower market and a popular cut flower in Holland, Germany and USA [2,3] In India, commercial production of gerberas is centred around Pune and Bangalore, parts of Sikkim, Nagaland, Meghalaya and Uttarakhand, from where flowers are being sent to local and international market.

The plant growth and development as well as flower quality is significantly influenced by climatic factors as well as nutrient management. Due to imbalanced use of macro and micro nutrients, the plant growth, development and quality of flower are directly affected. Therefore, balanced nutrient application is necessary for healthy plant growth and production of quality flowers. Micronutrients play vital roles in the growth and development of plants, due to their stimulatory and catalytic effects on metabolic processes and ultimately on flower yield and quality [4]. These are essential for crop growth and development. Enrichment of crop with micronutrients, especially zinc, iron and magnesium may prove effective in regulating flowering in crops [5] and aid in flower production.

Zinc is necessary for the synthesis of auxin IAA and for carbohydrate metabolism, protein synthesis [6]. Iron is a key element in different oxidation-reduction reactions of respiration and photosynthesis [7] and magnesium is involved in chlorophyll synthesis, enzyme activation and protein synthesis. Micronutrients are to be necessarily taken up by the plants from soil or supplemented through foliar application for good growth and yield of crops. Foliar application of nutrients is gaining more importance in fertilization of various field and floricultural crops,

in many countries. Foliar fertilization technique may also be a good alternative to the conventional soil application to avoid the loss of fertilizers by leaching. It is recommended by several investigators as an alternative fertilization method to improve the growth and flowering of gerbera. Keeping these facts in view, this study evaluated the effect of foliar application of Zn, Fe and Mg on growth, flowering and yield of gerbera under protected condition.

## 2. MATERIALS AND METHODS

An experiment was carried out on "Effect of foliar application of micronutrients on growth, yield and flower quality of gerbera (*Gerbera jamesonii*) under naturally ventilated polyhouse conditions in Prayagraj" at Horticulture Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj, during 2022-2023.

The experiment was laid out in Randomized Block design with ten treatments and replicated thrice that is, control (T0), ZnSO<sub>4</sub> @ 0.2% (T1), ZnSO<sub>4</sub> @ 0.4% (T2), ZnSO<sub>4</sub> @ 0.6% (T3), FeSO<sub>4</sub> @ 0.2% (T4), FeSO<sub>4</sub> @ 0.4% (T5), FeSO<sub>4</sub> @ 0.6% (T6), MgSO<sub>4</sub> @ 0.2% (T7), MgSO<sub>4</sub> @ 0.4% (T8) and MgSO<sub>4</sub> @ 0.9% (T9). Healthy tissue cultured plants having 4-6 leaves were planted on flat beds. The planting material were procured from Sheel- Biotech, New-Delhi. They were planted at a spacing of 30 cm x 30 cm. As per treatments the micronutrients were sprayed after one month of planting and subsequently at one-month intervals up to 4 months. Four plants from each replication of a cultivar were used for recording observation. The recommended cultural practices were followed for raising the successful crop. Data on growth, flowering, yield and quality characters was recorded up to 4 months after one month of planting.

## 3. RESULTS AND DISCUSSION

### 3.1 Vegetative Parameters

The effect of levels of micronutrients on vegetative parameters of Gerbera are shown in Table 1. From table 1, the application of different levels of Zn, Fe and Mg had produced a significant effect on growth parameters, viz. plant height, plant spread, number of leaves.

Significantly taller plants (24.08 cm) were observed with treatment T2 (Zn@0.4%) followed by T5 (Fe@0.4%, 22.63 cm) whereas, significantly shorter plants were reported in T0 (Control, 17.69 cm). The increase in plant height with application of micronutrient Zn might due to its role in the synthesis of tryptophan which is a precursor of auxin (IAA). Similarly, Fe acts as an important catalyst in enzymatic reactions of metabolism and would have helped in larger biosynthesis of photo assimilates thereby enhancing the growth of plants. Similar results were reported by in Balakrishnan et al. [8] in marigold and Pal et al., [9] in gerbera. Significantly wider plant spread (m<sup>2</sup>) (37.68 cm) was observed with treatment T2 (Zn@0.4%) followed by T5 (Fe@0.4%, 35.80 cm) whereas, significantly lesser plant spread (m<sup>2</sup>) was reported in T0 (Control, 23.62 cm). The plant spread significantly influenced with application of micronutrient ferrous sulphate is an essential component of several dehydrogenase, proteinase, peptidase and promotes growth hormones and closely associated with plant growth, all these factors contribute to cell multiplication, cell division and cell differentiation resulting in increased photosynthesis and translocation of food material which enhanced the plant spread and zinc sulphate could have attributed to improved root system of plants resulting in absorption of water and nutrients and its utilization. Similar results were also obtained by Kakade et al., [10] in China aster, Balakrishnan [8] in marigold. Significantly more leaves (15.08) were recorded in treatment (Zn@0.4%) followed by T5 (Fe@0.4%, 11.17) and significantly a smaller number of leaves in T0 (Control, 7.00). Zinc (Zn) and iron (Fe) are

essential micronutrients for plant growth and development including photosynthesis, enzyme activation, and hormone synthesis. Adequate levels of these micronutrients help in promoting leaf growth and increasing leaf number. Similar results were reported by Khan [11]. The statistically analysed data was presented in Table 1.

### 3.2 Floral Parameters

The effect of levels of micronutrients on floral parameters of Gerbera are shown in Table 2. From Table 2, the application of different levels of Zn, Fe and Mg exerted a significant influence on flowering and yield parameters, viz. days for first bud initiation, number of flowers/plant, stalk length and vase life. The review of data presented in Table 2 indicated that significantly lesser days (73.25) were taken for first bud initiation in treatment (Zn@0.4%) followed by T5(Fe@0.4%, 85.58) while significantly more number of days taken to first bud initiation T0-Control (106.67). Micronutrients like zinc and iron aids in storage of more carbohydrates through photosynthesis, which may be enhanced to earlier flower bud initiation. Similar results were reported by Jadhav et al. [12], Sujatha et al. [3] and Pimple et al. [13]. Significantly more flowers per plant (10) was found with plants treated with treatment (Zn@0.4%) followed by T5(Fe@0.4%, (9), whereas significantly lower flowers were recorded in T0-Control (6). Application of iron and zinc enhances vegetative growth and healthy green leaves which in turn resulted in higher assimilate synthesis which may increase the flower production and ultimately flower yield. Similar results were obtained by

**Table 1. Effect of different levels of micronutrients on vegetative parameters of gerbera**

Treatment Symbol	Treatment Combination	Plant height(cm)	Plant spread (cm)	Number of leaves per plant
T <sub>0</sub>	Control	17.69	23.62	7.00
T <sub>1</sub>	Zn@0.2%	21.43	33.61	8.67
T <sub>2</sub>	Zn@0.4%	24.08	37.68	15.08
T <sub>3</sub>	Zn@0.6%	22.55	35.38	9.67
T <sub>4</sub>	Fe@0.2%	22.37	32.43	7.17
T <sub>5</sub>	Fe@0.4%	22.63	35.80	9.58
T <sub>6</sub>	Fe@0.6%	21.80	35.70	11.17
T <sub>7</sub>	Mg@0.2%	20.39	34.86	7.83
T <sub>8</sub>	Mg@0.4%	18.86	34.50	7.67
T <sub>9</sub>	Mg@0.6%	18.50	26.99	7.25
<b>F-Test</b>		<b>S</b>	<b>S</b>	<b>S</b>
S.Ed(±)		0.43	0.85	0.55
C.V.		2.52	3.15	7.47
CD 0.05		0.90	1.79	1.16

**Table 2. Effect of different levels of micronutrients on floral parameters of gerbera**

Treatment Symbol	Treatment Combination	Days to first bud emergence (Earliness)	Number of Flower/Plant	Stalk length(cm)	Vase life
T <sub>0</sub>	Control	<b>106.67</b>	8	<b>47.67</b>	<b>5.67</b>
T <sub>1</sub>	Zn@0.2%	95.17	9	49.25	6.86
T <sub>2</sub>	Zn@0.4%	<b>73.25</b>	<b>10</b>	<b>56.50</b>	6.33
T <sub>3</sub>	Zn@0.6%	89.08	8	49.08	<b>8.75</b>
T <sub>4</sub>	Fe@0.2%	87.00	9	49.17	7.22
T <sub>5</sub>	Fe@0.4%	85.33	<b>9</b>	<b>54.17</b>	<b>7.68</b>
T <sub>6</sub>	Fe@0.6%	<b>85.58</b>	8	50.50	6.89
T <sub>7</sub>	Mg@0.2%	93.00	6	48.42	6.44
T <sub>8</sub>	Mg@0.4%	100.42	6	48.58	6.69
T <sub>9</sub>	Mg@0.6%	98.42	<b>6</b>	49.67	6.00
<b>F-Test</b>		<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>
S.Ed(±)		3.71	0.43	0.93	0.49
C.V.		4.97	6.74	2.26	8.76
CD 0.05		7.80	0.90	1.95	1.03

Anuprita et al. [14] in gerbera and Pal et al. [9]. The highest stalk length (56.50 cm) was observed with treatment (Zn@0.4%) followed by T<sub>5</sub> (Fe@0.4%, 54.17 cm) while lowest was reported in T<sub>7</sub>(Mg@0.2%, 47.67 cm). The increment in the stalk length might be due to enhanced cell division and cell enlargement. Similar results were reported by Khan [11], Sujatha et al. [3] and Gurav et al. [15] in gerbera. More vase life (8.75) was observed with treatment (Zn@0.4%) followed by T<sub>5</sub>(Fe@0.4%, 7.68) whereas less vase life was reported in T<sub>0</sub>(Control, 5.67). The positive impact of micronutrients like zinc, iron and magnesium might be due to the ability of these nutrients in activating several enzymes and their involvement in chlorophyll synthesis and various physiological activities ultimately enhance the vase life. Similar results were reported by Jadhav et al. [12] and Pal et al. [9].

#### 4. CONCLUSION

It is concluded from the present investigation of 10 treatments under study that there was significant variation in all the parameters investigated. The treatment T<sub>2</sub>-Zn@0.4% reported significantly better performance in parameters like plant height, plant spread, earliness for first bud initiation, number of flowers per plant, stalk length, vase life.

#### ACKNOWLEDGEMENT

I express my gratitude to my Advisor Dr. Urfi Fatmi for constant support, guidance and valuable suggestions for improving the quality of

this research work and to all the faculty members of Department of Horticulture, SHUATS, Prayagraj, Uttar Pradesh for providing all necessary facilities, their cooperation, encouragement and support.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Sindhu SS. Popularizing flower cultivation for export. Indian Horticulture. 2006; 48(2):41-46.
2. Choudhary A, Prasad CA. Effect of potassium and calcium concentrations in the nutrient solution on growth and nutrient uptake of rose plants. International Journal of Science Research. 2000;69(4):606-608.
3. Sujatha K, Narayana GJV, Khan MM. Effect of different fertigation levels on gerbera under low-cost greenhouse. Journal of Ornamental Horticulture. 2002; 5(1):54-9.
4. Khosa SS, Younis A, Yameen S, Riaj A. Effect of foliar application of micro nutrient on growth and flowering of gerbera (*Gerbera jamesonii*). American-BIBLIOGRAPHY 78 Eurasian Journal of Agriculture and Environmental Sciences. 2011;11(5):736-757.
5. Pratap M, Reddy SA, Reddy YN. Effect of foliar application of FeSO<sub>4</sub> and ZnSO<sub>4</sub> on flower production and anthocyanin content

- of *Gladiolus* spikes. Journal of Ornamental Horticulture. 2005;7(2):159-163.
6. Shukla AK, Dwivedi BS, Singh VK, Gill MS. Macro role of micronutrients. Indian Journal of Fertilizers. 2009;5(5):11– 2, 15–8, 21–4 and 27–30.
  7. Reddy YT, Reddi SGH. Principles of Agronomy, pp 204–6. Kalyani Publishers, Ludhiana; 2002.
  8. Balakrishnan V, Jawaharlal M, Senthil Kumar T, Ganga M. Response of micronutrients on flowering, yield and xanthophyll content in African marigold (*Tagetes erecta* Linn.). Journal of Ornamental Horticulture. 2007;10(3):153-156.
  9. Pal S, Barad AV, Singh AK, Khadda BS, Kumar D. Effect of foliar application of Fe and Zn on growth, flowering and yield of gerbera (*Gerbera jamesonii*) under protected condition. Indian Journal of Agricultural Sciences. 2016;86(3):394-398.
  10. Kakade DK, Rajput SG, Joshi KL. Effect of foliar application of Fe and Zn on growth, flowering and Yield of China aster (*Callistepus chinensis* Linnaeus). Asian Journal of Horticulture. 2009;4(1):138-140.
  11. Khan. Effect of micronutrients on dahlia. Journal of Ornamental Horticulture. 2000; 3(2):122–3.
  12. Jadhav AH, Dalal SR, Shinde RD, Deshmukh RP. Effect of micronutrients on growth and flower production of gerbera under polyhouse conditions. Advances in Plant Sciences. 2005;18(2):755-758.
  13. Pimple AG, Dalal SR, Nandre DR, Ghawade SM. Yield and quality of gerbera influenced by nitrogen and phosphorus levels under polyhouse conditions. International Journal of Agricultural Science. 2006;2(2):320–1.
  14. Anuprita H, Jadhav SR, Dalal RD, Deshmukh RP. Effect of micronutrients on growth and flower production of gerbera under poly house conditions. Advances in Plant Science. 2005;18(2):755–8.
  15. Gurav SB, Katwate SM, Singh BR, Sable RN, Kakade DS, Dhane AV. Effect of nutritional levels on yield and quality of gerbera. Journal of Ornamental Horticulture. 2004;7(3-4):226–9.

© 2023 Mounika and Fatmi; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/104662>