

Asian Journal of Pediatric Research

Volume 14, Issue 3, Page 27-36, 2024; Article no.AJPR.112024 ISSN: 2582-2950

Family Centered Care: The Perspective of Nurses in the Neonatal Intensive Care Unit of Tamale Teaching Hospital, Ghana: A Qualitative Exploration

Victoria Awalenkak Agwiah ^{a,b} and Keren-Happuch Twumasiwaa Boateng ^{b,c*}

^a Department of Paediatrics and Child Health, Tamale Teaching Hospital, Tamale, Ghana.
 ^b Midwifery and Paediatrics Division, Ghana College of Nurses and Midwives, Accra, Ghana.
 ^c Department of Midwifery and Women's Health, School of Nursing and Midwifery, University for Development Studies, Tamale, Ghana.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJPR/2024/v14i3331

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

https://www.sdiarticle5.com/review-history/112024

Original Research Article

Received: 05/12/2023 Accepted: 09/02/2024 Published: 07/03/2024

ABSTRACT

Background: Family-Centered Care is a care approach in which healthcare professionals collaborate with the patient's family to plan and make decisions regarding the patient's treatment. Family-Centered Care is widely acknowledged and documented as the optimal care approach for neonatal practice. Despite the growing use of this method, there is a lack of research on the perspectives of healthcare professionals and carers regarding Family-Centered Care. The study aimed to evaluate the viewpoints of nurses regarding FCC in the newborn critical care unit at the Tamale Teaching Hospital.

Methodology: The study utilised an exploratory descriptive design employing a qualitative technique, in which focus group talks were held among participants. The material was transcribed and processed via thematic and content analysis techniques.

*Corresponding author: Email: khappuch89@gmail.com;

Findings: Approximately majority of the participants demonstrated awareness of the idea of Family-Centered Care, which they elaborated upon through two main themes namely provision of care for both the patient and their family, and Involvement of relatives in the care of the patient. **Recommendations:** Majority of participants expressed that Family-Centered Care was not practiced in their unit. Nevertheless, all participants unanimously concurred that the approach had the potential to yield positive results in terms of patient care outcomes.

Keywords: Family centered care; neonatal intensive care unit; neonates; nurse perspectives; Tamale Teaching Hospital

1. INTRODUCTION

Family Centered Care (FCC) is the model of care where health professionals plan and make decision together with the patients' family in care of their patient [1]. A focus on improved new-born care practices is critical in reducing under-five mortality across settings [2]. Such focus is essential because an estimated 5.9 million children die before their fifth birthday worldwide [3]. In Sub-Sahara Africa alone, an estimated three (3) million deaths among one (1) month olds occurred in 2015 with one (1) death occurring in every twelve (12) under-five (UN MDG, 2015; World Bank, 2015). The Ghana Maternal and Health Survey reported a childhood mortality of twenty-five (25) deaths per 1000 live births in the country [4]. The neonatal period, which spans the first 28 days of life, is a critical and vulnerable phase for the survival of neonates. Neonatal mortality accounts for an average of 60% of all infant mortality in Ghana with twenty-nine (29) deaths per 1,000 live births Consequently, neonatal mortality become a very important component of infant and child mortality and requires very urgent attention (Ghana National Newborn Health Strategy, 2014).

The Neonatal Intensive Care Unit is a setting specially equipped with trained staff and resources to care for sick babies especially those born premature for effective care. hospitalizations present a challenge to parents as they are usually restricted entry to the NICU during the care of their premature babies due to limited infrastructure or hospital protocols [6]. The situation often excludes parents from important discussions and are left in the dark about the management of their Children's condition [7] which result in further aggravation of the already stressed parents leading to anxiety and despair [8,9]. Also, early separation of premature babies from their mothers in the postnatal period have been reported to be stressful to babies [10] and impair their emotional, cognitive and psychomotor development [11].

Similarly, inadequate bonding between mothers and babies results in improper initiation of breastfeeding [12]. The moment nurses and doctors start to take care of babies in the hospital, parents are reported to lose their sense of control and this birthed the concept of FCC where nurses share the care of children with their families [13]. Several intervention programs have since been designed across many settings including Ghana to care for babies and their parents including Kangaroo Mother Care (KMC), FCC and Skin-to-skin care [4]. Among these intervention programs, FCC is considered the primary model of care for premature babies and their families in the NICU [14].

In the FCC model of health care, family members play a role in all aspects of health care for their babies [15]. The Institute for Patient and FCC (IPFCC) defines the concept of FCC as an approach to organize, deliver, and evaluate the health care that is grounded in mutual valuable collaboration amid the health care providers, patients and their families [16]. Also, the Institute of Medicine (IOM) defines FCC as the care that is respectful and responsive to individual patient preference, needs and values (IOM, 2010).

In neonatal intensive care (NIC), the concept of FCC refers to the recognition given to families as being integral to the healthcare team [17]. In FCC, healthcare is designed to provide care for both the family and the child, rather than solely focusing on the individual child. For FCC to be fully effective, parents should be allowed to enter the NICU without time restriction [18 19,20].

The FCC approach of care in the NICU has been found to be beneficial to both premature babies and parents [21]. For example, FCC was found to have decreased the length of stay of patients in NICU in the Netherlands [3]. Similarly, FCC

through skin-to-skin care have been found to improve infants cardiac and respiratory system, stabilize temperature, organised sleep, breastfeeding and neuro-developmental outcome [22].

Also, the benefits of FCC to parents of premature babies have been reported in several studies resulting in reduced stress level among parents in Germany [23], increase parents' satisfaction care in the Netherlands [2] empowerment of parents to have self-confidence and competences in assuming their roles after discharge in the USA [24]. Furthermore, [25] found a positive association of FCC with improvements in efficient use of services, health satisfaction, access to communication, systems of care and family functioning.

2. METHODS

2.1 Study Design

An exploratory descriptive design, which fits within a qualitative approach, was employed to explore the perspectives of nurses on FCC in Tamale Teaching Hospital. This design enabled investigator have to an in-depth understanding of the phenomenon [26]. This design was chosen in this study because the researcher is interested in understanding the knowledge and attitudes towards Family-Centered Care among Neonatal Intensive Care Unit nurses. Unlike quantitative study, qualitative study's major focus is to collect quality data for the purposes of describing and understanding the real meaning of the research problem [27].

2.2 Setting

Tamale Teaching Hospital was the study center and was set up in 1974 and was formerly the Northern Regional Hospital. Ιt was upgraded in 2005 to a teaching hospital. The hospital is the only teaching hospital in the Northern part of Ghana. The hospital provides health service and receives referrals from the regional and districts hospitals in the region and the surrounding regions. The hospital also provides clinical training serves to the University for Development Studies, Medical School and School of Nursing and Midwifery. Neonatal Intensive Care Unit of the TTH was the site for the study. The unit has a bed capacity of 70 and staff strength of 42.

2.3 Target Population

The target population for this study were all nurses working in the neonatal intensive care unit of the Tamale Teaching Hospital for at least three months.

2.3.1 Inclusion criteria

The study included all nurses working in the NICU for at least three months.

2.3.2 Exclusion criteria

The study excluded all nurses working in NICU whose children were on admission in the same unit

Sampling Technique and Size: Purposive sampling technique was used to engage eight (8) participants, data saturation was reached given rise to a sample size of eight (8) nurses. Data saturation was at the point where the researcher no longer gets new insights from the data (Creswell, 2014). This technique was used when the participants have the requisite knowledge on the topic under study [28]. As the researcher will recruit participants that will give relevant data until data saturation level is achieved.

2.4 Data Collection Procedure

The researcher developed the interview guide based on the objectives of the study. A tape recorder was used with the permission of participants to record the interview for adequate interpretation. The researcher also used pens and jotters to take notes of important non-verbal clues to confirm and support verbal responses. The data was enriched with probing questions. Before the start of the interview, each participant was guided to complete the participants' background information sheet. This helped the researcher to understand each participant's background and to also facilitate participants easy transit into the sharing of their experiences in the interview. The interview was audiotaped with the permission from the participants. The participants were asked questions on their experience on FCC. During each interview session, responses of participants were robed or redirected to focus their responses on the objectives of the study. The non-verbal cues such as change of voice tone, eyes tearing, and facial expressions were also noted on the field notes. Each interview lasted approximately 45-50 minutes [29] notes and audio records were

labelled and then stored in the researcher personal drawer under lock and key. The data will be kept for 5 years for the purpose of audit trail.

3. METHODOLOGICAL RIGOUR

Trustworthiness: This research employed the trustworthiness criteria recommended in the works of [30] to ensure methodological rigour. This included credibility, transferability, dependability, and conformability [31,32].

Credibility involves processes to ensure that the participants can confirm results [31,33,34]. Data triangulation and prolonged engagement in the subject matter ensured this [35].

Dependability refers to ensuring acceptable procedures and processes have been followed to establish the study's trustworthiness [36]. To achieve this, there was a provision of a detailed report of the processes involved in the study, which included (1) a description of the research design and implementation, (2) a detailed explanation of the data gathering process and (3) explaining what occurred in the field. An experienced qualitative researcher also did independent co-coding of data compared for analysis. Triangulation data determined consistency and stability across the various data sources and methods employed in the study. Dependability is comparable with reliability [31].

Confirmability involves verifying that the obtained data is accurate and relevant, and that it supports the results and recommendations made in the study. This was met through triangulation of the methods (a combination of field notes and interviews) and keeping of audit trails, as well as member checking [32] by taking the final report back to the participants to determine whether they felt that it was an accurate representation of their input (Creswell and Plano-Clark, 2011). It is comparable with objectivity or neutrality [31].

Transferability refers to the applicability of the study. This enables other researchers to use the results on similar participants in a similar context. This was addressed by providing detailed descriptions as presented by the words and phrases used by participants; this comprised sufficient contextual information about the fieldwork to enable the reader to make such a transfer. Also presented was a detailed description of the study setting, the respondents,

and the methods involved. Transferability is comparable with external validity [31].

3.1 Study Findings

The study primarily focused on nurses' perspectives on FCC. The major themes emerged were Knowledge of FCC among nurses and Practice of FCC at the NICU. The subjects were introduced and exact quotations were employed to support the assertions.

3.2 Participants Demographics

This outlines the background characteristics of all 8 participants who participated in the focus group discussion. Out of the participants, 4 individuals (50%) were between 25-30 years old, 2 participants (25%) were aged 31-35 years, and 2 (25%) were aged 36-40 years. Out of the participants, 5 (60%) were ladies and 3 (40%) were males. Out of the participants, 5 individuals (60%) were married and 3 individuals (40%) were single. The majority of participants, 5 out of 8 (60%), were Christians, while 3 out of 8 (40%) were Muslims. Out of the participants, 5 (60%) had a degree and 3 (40%) had a diploma as their highest educational qualification.

3.3 Knowledge of Family Centered Care among Nurses

During the interaction, 6 (70%) out of the 8 respondents said they ever heard of FCC and only 2(30%) of them said she had never heard about it. In their responses to the second item (What is FCC) two (2) main themes emerged from the data were FCC as care for both patient and family; FCC as care of patient by involving patient relatives"

3.3.1 Family/Relatives participation in the care process

Participants in this study held varied explanations as to what constitute FCC even though they were familiar with the concept. R3 describes FCC in the following passage:

Like the colleague said even though I have not really heard it mentioned, I do understand what the concept is about. But if I understand it is the direct opposite to patient centered care. So, whereas patient centered care is about eh eh... taking care of only the patient, FCC deals with taking care of the baby and caregivers as well.(R3)

In a similar way, a participant described the concept of FCC and had this to say:

I know it is about offering holistic care. It is a case where.....probably not only the baby is cared for by the health team but all other persons who matter(R4)

The view of another participant was not different from what the others had said about FCC. The participant, also argues:

it is a form of care given to both patients and their families. It means that we don't only concentrate on the patient but also the... those who brought the patient from home.(R8)

All the participants had similar views regarding the fact they appreciate the FCC as being practiced in the unit.

3.3.2 FCC as care of patient by involving patient relative

From the focus group discussion, some participants argued that FCC was care of patient by involving the family. In his attempt to explain the concept of Family-Centered Care, R1, had to say in the following passage:

FCC...FCC. I have not really heard about it but em em, what I can say about it is that it is the care given to family let me say care centered on the whole family. (pause) like nuclear family of a sort where we involve them in care. (R1)

This explanation was reinforced by the explanation given by:

A holistic way of nursing where both the patient and relatives of the patient are nursed. In this case there might be need for some assessment to be done on the patient including involving the care givers in planning care.(R7)

It was established in this study that the participants were not too sure of the meaning of the concept of FCC. This was noted in the words of participant:

care of patients while meeting the needs of families of the Yes, I have heard about it but now and don't really hear about it. I'm not sure what exactly it is about. But if it is like the way they are describing it (silence) then well... about taking patient (**R5**)

However, there was one (1) participant, who simply said "no idea" admitting he did not know about the concept of FCC.R1

3.4 Practice of FCC at the Neonatal Intensive Care Unit

On the practice of FCC, 7 out of the 8 participants indicated that FCC was not practiced in the ward. Responses to the questions under the practice objective produce two (2) central themes; parents are not involved in the care of the sick" "parents are partially involved in the care of the sick.

3.4.1 Parents are not involved in the care of the sick

This study found majority of the participant nurses stating that FCC was not practiced in the neonatal intensive care unit. Participant did not hesitate to state his position in the passage below:

Oh no no, we usually just take care of the baby all alone as health workers. We don't allow that all. That is why we have enough space for their mothers to always stay waiting for their babies. (R1)

One participant, who probably abhorred the practice of FCC had this to say:

Not here. You know (pause), we do not have a policy that says relatives should assist in care of babies. So, we accordingly do not do that. we do the assessment of the baby's situation and we carry out the care all alone. We decide when they should take medication, when to check the vital signs of the child even decide the topics on which we educate them. (**R6**)

3.4.2 Parents are partially involved in the care of the sick

However, the nurses found the need to involve them on a number of occasions. Participant in her expression had this to say:

We only call them when the child is hungry and crying for food. What possibly can they do to help apart from feeding the baby? We don't let them take part, yaa. (R2)

Some participants also blamed the exclusion of caregivers in the care of the sick on physician activity such as conducting ward rounds. Participant was explicit in his statement:

family members are usually sacked especially when doc comes for review. They are not really allowed to participate in the care at all. (R3)

The participants also argued that caregivers are involved when it came to purchasing drug and other services:

I don't think we do that here in this ward. We make them sit comfortably while we reassure them of competent care. But when it comes to buying of drugs, we call them to go buy some for us. But as for actual nursing care relatives are not taking part in this unit.(R5)

involvement views as expressed by participant R4 was not too different that of the others:. For instance, if you ask the mother of a child to hold down a baby so that you the care.(....... I will say yes and I say so because any form or level of involvement is The nurse will cannulate the baby, what is that? On that score, I will say they take part in patient's R4)

5. DISCUSSION

This presents the detailed discussion of the major findings of this study in association with existing relevant literature pertinent to FCC. The discussion is presented according to the major themes and subthemes which were based on the objectives of the study.

5.1 Knowledge of FCC among Nurses

This study found that about three-quarters of the participants were aware of the concept of FCC. This study finding was reflected in the responses of the participants as they shared their views on what FCC was. However, participant's knowledge towards FCC varied. This study findings concurs with what has been reported by various authors [37]. This finding could have been influenced by exposure to the various explanations or definitions of the concept of FCC as documented by [38], [39] and [40]. This reality

of nurses giving varied views of FCC could be contributing to the ongoing debate that nurses misunderstand the concept of FCC as argued by [38]. Comparing the definitions given by the participants to those articulated in literature by various 40, some differences in meaning were noted. Notwithstanding, participants covered only one of the two key components of the concept. It was also noted that, participants understood the concept as having more to do with the care of the care giver as well as the client. Caregiver involvement which is a key component of FCC not captured frequently. Again, this study revealed that participants did not know the typical definition of the concept of Family-Centered Care. They provided uncertain responses indicating a lack of understanding of the concept. However, individuals provided replies that closely aligned with the subject being studied. Their knowledge of this concept could have been influenced by their exposure to similar concepts like patient centered care.

The findings in this study are similar to the findings of [41] study on FCC in Newborn Unit: Nursing perspective. They found that the application of FCC is wrongly understood, and portrays the unpreparedness of professionals to deal with the Family process of hospitalised children.

5.2 Practice of FCC at the Neonatal Intensive Care Unit

On the practice of FCC, this study found that more than half of the respondents saying that FCC is not implemented in the unit. The participants were very straight forward with their answers and went ahead to specify what they meant by not practicing FCC. For instance, they likened FCC with care giver involvement during the care process and said they did not involve caregivers during care. This finding was consistent with that of [42] and [18] and [43] when they suggested the implementation of FCC was problematic in health facilities.

The study also found that nurses occasionally involved care givers in the care of patients. This was noted when the participants argued that caregivers are asked to purchase drugs and also feed their babies. This study revealed a number of factors that accounted for nurses' inability to implement FCC. The factors cited include workload, burden of documentation, confused state of relatives and physician ward rounds fear of relatives interfering with clinical duties. This

finding was consistent with what has been reported by [44] who argued that such factors as unit culture, staff resistance, lack of space and time, clinicians being uncomfortable with family being present, and uncertainty about the benefits of FCC were barriers to the practice of FCC. This study also revealed that parents or care givers needs may be attended to by Nurses but they often did so when caregivers were sick [45-47].

It was also demonstrated that, respondents did not carry out home visiting or follow-up visits. However, the study found commitment to follow-up review where the patient and relatives are expected to come back for further assessment and treatment. This finding was in line with the placement or recruitment policy of the Nurses in this ward. This was probably so because all the nurses were registered nurses who are trained to offer services in the wards. It probably was on this basis that one of the participants reported that home visiting was not feasible.

The discovery in this study was consistent to with the findings of [1] study on FCC from the perspective of Nurses in the Neonatal Intensive Care Unit. They found lack of material resources, shortage of staff, creating and updating protocols, training as challenges hindering FCC.

6. CONCLUSION

Family-Centered Care is a key component of care for neonates at neonatal intensive care units. Its benefits are so keen in the care of neonates in reducing neonatal mortality and morbidity, yet it lacks proper stance in the care of neonates in the Tamale Teaching Hospital NICU. The nurses reported some potential benefits and challenges of adopting FCC approach in the delivery of care. This study highlighted the need for FCC to be used as an approach to planning, delivery, and evaluation of health care that is grounded in mutual beneficial partnership among health care providers, patients and families.

CONSENT AND ETHICAL APPROVAL

An introductory letter from the research committee of the Ghana College of Nurses and Midwives was sent to the Ethics Review Committee of the Navrongo Health Research Centre for ethical clearance before data collection. The researcher also sought permission from management of the Tamale

Teaching Hospital before the data collection. The participant's consent was sought by them agreeing and filling a consent form to participate in the study. The participants were given the opportunity to either participate or opt out from the study at any level they desired. The study participants were also educated that consent covers tape recording of the interview, notes taking of observations that cannot be recorded by the tape, description and reporting of findings. The study participants were assured of confidentiality of all the information given during the interview. As such, tape recorder and notebook were constantly under lock and key when not in use and transcript and computed information were pass-worded. Participants were informed not to use their names or any form of identification. The participants were guaranteed of no harm throughout the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Silva TRG, Manzo BF, de Faria Fioreti FCC, Silva PM. Family-centered care from the perspective of nurses in the Neonatal Intensive Care Unit. Revista da Rede de Enfermagem do Nordeste. 2016;17(5):643-650.
- Segers E, Ockhuijsen H, Baarendse P, Eerden IV, van den Hoogen A. The impact of family centred care interventions in a neonatal or paediatric intensive care unit on parents' satisfaction and length of stay: A systematic review. Intensive and Critical Care Nursing. 2018;1-8.
 - DOI:10.1016/j.iccn.2018.08.008
- The United Nations. The millennium development goals report; 2015. Accessed 01 Mar 2024.
- Ghana Statistical Service, Macro International. Ghana demographic and health survey 2018. Accra. Ghana: Ghana Statistical Service, Ghana Health Service. and **ICF** Macro: 2019.
- Ghana Statistical Service, United Nations Children's Fund. Ghana Multiple Indicator Cluster Survey (MICS) 2011, Ref. GHA_2011_MICS_v01_M. Available:http://www.statsghana.gov.gh/na da/index.php/catalog/52.

- 6. De Vonderweid U, Leonessa M. Family centered neonatal care. Early Human Development. 2009;85:37-38. DOI:10.1016/i.earlhumdev.2009.08.009
- 7. Barry M, Edgman-Levitan S. Shared decision making The pinnacle of patient-centered care. The New England

Journal of Medicine. 2012;780-781.

- 8. Franck LS, McNulty A, Alderdice F. The perinatal-neonatal care journey for parents of preterm infants. The Journal of Perinatal and Neonatal Nursing. 2017;31(3):244-
- Minton C, Batten L, Huntington A. A multicase study of prolonged critical illness in the intensive care unit: Families' experiences. Intensive Crit Care Nurs; 2018.
 - DOI:10.1016/j.iccn.2018.08.010
- Császár-Nagy N, Bókkon I. Mothernewborn separation at birth in hospitals: A possible risk for neurodevelopmental disorders? Neuroscience and Biobehavioral Reviews. 2018;84:337-351. DOI:https://doi.org/10.1016/j.neubiorev.201 7.08.013
- Ahlqvist-Björkroth S, Boukydis Z, Axelin 11. AM, Lehtonen L. Close collaboration with parents[™] intervention to improve parents' psychological well-being and child development: Description of the intervention and study protocol. Behavioural Brain Research, 2017;325;303-310.
- 12. Nyqvist KH, Häggkvist AP, Hansen MN, Kylberg E, Frandsen AL, Maastrup R, Haiek LN. Expansion of the baby-friendly hospital initiative ten steps to successful breastfeeding into neonatal intensive care: expert group recommendations. Journal of Human Lactation. 2013;29(3):300-309.
- 13. Tufekci FG, Kliniklerinde C, Bakim AM, Uygulamalarinin. The examination of family centered care practices in pediatric clinics. Journal of Anatolia Nursing and Health Sciences. 2015;18(2):131-139.
- De Bernardo G, Svelto M, Giordano M, Sordino D, Riccitelli M. Supporting parents in taking care of their infants admitted to a neonatal intensive care unit: a prospective cohort pilot study. Ital J Pediatr. 2017;43(1):36. DOI:10.1186/s13052-017-0352-1
- 15. Roets L, Rowe-Rowe N, Nel R. Family centred care in the paediatric intensive care unit. Journal of nursing management. 2012;20(5):624-630.

- 16. Institute for patient and family-centered care, "frequently asked questions,"; 2014. Available: http://www.ipfcc.org/fag.html.
- Boykova M, Kenner C. Family as the center of care. Newborn and Infant Nursing Reviews. 2016;16: 40.
 DOI:10.1053/j.nainr.2016.03.003
- Finlayson K, Dixon A, Smith C, Dykes F, Flacking R. Mothers' perceptions of family centred care in neonatal intensive care units. Sexual and Reproductive Healthcare. 2014;5:119–124. DOI:10.1016/j.srhc.2014.06.003
- Latour J, Van Goudoever J, Schuurman BE, Albers M, Van Dam N, Dullaart E, Hazelzet JA. A qualitative study exploring the experiences of parents of children admitted to seven Dutch pediatric intensive care units. Intensive Care Medicine. 2011;37:319–325. DOI:10.1007/s00134-010-2074-3)
- 20. Thiele N, Knierim I, Mader S. Parents as partners in care: Seven guiding principles to ease the collaboration. Newborn and Infant Nursing Reviews. 2016;16:66–68. DOI:10.1053/j.nainr.2016.03.012
- 21. Silva TRG, Manzo BF, Fioreti FCCF, Silva PM. Family-centered care from the perspective of nurses in the neonatal intensive care unit. Rev RENE. 2015;17(5):643-50.
- Skene C, Gerrish K, Price F, Pilling E, Bayliss P, Gillespie S. Developing familycentred care in a neonatal intensive care unit: An action research study. Intensive and Critical Care Nursing; 2018. DOI:10.1016/j.iccn.2018.05.006
- 23. Enke C, Oliva Y, Hausmann A, Miedaner F, Roth B, Woopen C. Communicating with parents in neonatal intensive care units: The impact on parental stress. Patient Education and Counseling. 2017;100:710-719.
 - Available:
 - https://doi.org/10.1016/j.pec.2016.11.017
- 24. Fleck P. Connecting mothers and infants in the neonatal intensive care unit. Newborn and Infant Nursing Reviews. 2016;16:92–96.
 - DOI:10.1053/j.nainr.2016.03.007
- Kuhlthau KA, Bloom S, Van Cleave J, Knapp AA, Romm D, Klatka K, Perrin JM. Evidence for family centered care for children with special health care needs: A systematic review. Acad Pediatr. 2011;11(2):136-143. DOI:10.1016/j.acap.2010.12.014

- Mayan MJ. Essentials of qualitative inquiry. Walnut Creek, CA: Left Coast Press: 2009.
- 27. Creswell JW. Research design: Qualitative, quantitative and mixed methods approaches (4th ed.). Thousand Oaks, CA: Sage; 2014.
- 28. Khan M, Muhammad N, Ahmed M, Saeed F, Aman KS. Impact of activity-based teaching on students' academic achievements in physics at secondary level. Academic Research International. 2012;3: 146-156.
- Goyal P, Rahman Z, Kazmi A. Corporate sustainability performance and firm performance research: literature review and future research agenda. Management Decision. 2013;51:361-379. Available:https://doi.org/10.1108/00251741 311301867
- Lincoln YS, Guba EG. Naturalistic inquiry. newbury park, CA: Sage Publications; 1985.
- 31. Tobin GA, Begley CM. 'Methodological rigour within a qualitative framework', Journal of Advanced Nursing. 2004;4:388–396.
 DOI: 10.1111/j.1365-2648.2004.03207.
- 32. McBrien B. 'Evidence-based care: enhancing the rigour of a qualitative study.', British Journal of Nursing (Mark Allen Publishing). 2008;17(20): 1286–1289. DOI:10.12968/bjon.2008.17.20.31645.
- 33. Newman I, Lim J, Pineda F. 'Content validity using a mixed methods approach: Its application and development through the use of a table of specifications methodology. Journal of Mixed Methods Research. 2013;7(3):243–
 - DOI: 10.1177/1558689813476922.
- 34. Polit DF, Beck CT. Essentials of nursing research appraising evidence for nursing practice. Philadelphia: Lippincott Williams & Wilkins; 2013.
- 35. Butler Ashleigh E, Copnell B, Hall H. 'The impact of the social and physical environments on parent healthcare provider relationships when a child dies in PICU: Findings from a grounded theory study', Intensive & Critical Care Nursing. Elsevier Ltd. 2019;50:28–35.
 - DOI: 10.1016/j.iccn.2017.12.008
- 36. Brink H, Van der Walt C, Van Rensburg G. Fundamentals of research methodology for

- healthcare professionals. 3rd edn. Cape Town: Juta & Co: 2012.
- 37. Kuo Dennis Z, Amy J Houtrow, Polly Arango, Karen A Kuhlthau, Jeffrey M Simmons, John M Neff. Family-centered care: Current applications and future directions in pediatric health care. Maternal and Child Health Journal. 2011;162:297-305.
- 38. Shields L, Zhou H, Pratt J, Taylor M, Hunter J, Pascoe E. Family centred care for hospitalised children aged 0-12 years. Cochrane database of systematic reviews. 2006:10.
- 39. Gooding JS, Cooper LG, Blaine AI, Franck LS, Howse JL, Berns SD. Family support and Family Centered Care in the neonatal intensive care unit: origins, advances, impact. Semin Perinatol. 2011;35(1): 20-28.
 - DOI:10.1053/j.semperi.2010.10.004
- 40. Coyne I, Murphy M, Costello T, O'Neill C, Donnellan C. A survey of nurses' practices and perceptions of Family Centered Care in Ireland. Journal of Family Nursing. 2013;19(4):469-488.
- 41. Corrêa AR, Andrade ACD, Manzo BF, Couto DL, Duarte ED. The family centered care practices in newborn unit nursing perspective. Escola Anna Nery. 2015; 19(4):629-634.
- 42. Harrison TM. Family-centered paediatric nursing care: State of the science. Journal of Paediatric Nursing. 2010;25(5);335-343. DOI:https://doi.org/10.1016/j.pedn.2009.01
- 43. Kuo D, Houtrow A, Arango P, Kuhlthau K, Simmons J, Neff J. Family-centered care: Current applications and future directions in pediatric healthcare. Maternal and Child Health Journal. 2012;16:297-305.

 Available: https://doi.org/10.1007/s10995-011-0751-7
- 44. Kleinpell R, Heyland D, Lipman J, Sprung C, Levy M, Mer M, Curtis R. Patient and family engagement in the ICU: Report from the task force of the world federation of societies of intensive and critical care medicine. Journal of Critical Care. 2018;48:251-256.
 - DOI:10.1016/j.jcrc.2018.09.006
- Ministry of Health Ghana. Ghana National Newborn Health Strategy and Action Plan; 2014–2018.
- 46. Creswell JW, Plano-Clark VL. Designing and conducting mixed methods research.

2nd edn. London: Sage Publications Ltd; 2011.

47. Kuo DZ, Houtrow AJ, Arango P, Kuhlthau KA, Simmons JM, Neff JM. Family centered care: Current applications and

future directions in pediatric health care. Matern Child Health J. 2012;16(2):297-305.

DOI:10.1007/s10995-011-0751-7

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
https://www.sdiarticle5.com/review-history/112024