



Optimizing Growth Parameters of Bell Pepper (*Capsicum frutescens* L.) Through Comprehensive Nutrient Management on Farm of the Department of Horticulture, MGCGV, Chitrakoot, Madhya Pradesh, India

Amit Vikram Gangele^{a++*}, S. S. Singh^{b#}, Priya Singh^{c†},
Kanta Kumar Sahu^{d‡}, Devanshu Dixit^{d‡}
and Ankur Dwivedi^{c^}

^a Department of Horticulture, MGCGV Chitrakoot, Satna (M.P.), India.

^b Division of Horticulture, MGCGV Chitrakoot, Satna (M.P.), India.

^c MGCGV Chitrakoot, Satna (M.P.), India.

^d Department of Technology Transfer, MGCGV Chitrakoot, Satna (M.P.), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJSSPN/2024/v10i1251

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113938>

Original Research Article

Received: 02/01/2024

Accepted: 06/03/2024

Published: 21/03/2024

⁺⁺ Guest Lecturer;

[#] Assistant Professor;

[†] M. Sc. (Ag.) Horticulture;

[‡] Ph. D Scholar (Agril. Extension);

[^] Ph. D Scholar (Animal Husbandry & Dairying Science);

*Corresponding author: Email: gangeleamit2015@gmail.com;

ABSTRACT

The study was conducted at the Research Farm of the Department of Horticulture, MGCGV, Chitrakoot. Bell pepper, a popular vegetable crop, faces challenges related to growth and productivity. This research aimed to determine the most suitable treatment for enhancing the growth parameters of Bell pepper plants.

The experiment involved a comprehensive approach, combining organic nutrient management, bioenhancers, and chemical fertilizers. Various treatments were applied to evaluate their impact on the growth parameters of Bell pepper. These parameters included plant height, leaf area, flowering, fruit set, and yield. Additionally, soil health and nutrient dynamics were assessed to understand the long-term sustainability of the treatments.

Preliminary findings suggest that a balanced combination of organic nutrients, bioenhancers, and judicious use of chemical fertilizers significantly improved the growth parameters of Bell pepper. This approach not only promoted vigorous vegetative growth but also enhanced reproductive stages, leading to higher fruit set and ultimately increased yield. Moreover, the study provides insights into the potential for sustainable Bell pepper cultivation practices.

The outcomes of this research contribute to the development of effective strategies for optimizing the growth parameters of Bell pepper, thereby enhancing its overall productivity and sustainability in agriculture. Further investigation is recommended to validate and fine-tune these findings for broader adoption in Bell pepper cultivation.

Keywords: Growth parameters; bell pepper; nutrient management; commercial crops.

1. INTRODUCTION

Bell pepper (*Capsicum frutescens* L.) is an annual and day neutral plant belongs to Solanaceae family. It is a one of the most important vegetable crop grown in India as well as in the world, because of its nutritive value, flavor, colour and is considered as one of the major commercial crops of the world [1]. It is relatively non pungent or less pungent with thick flesh and is the world second most important vegetable crop after tomato. Sweet pepper has little energy value but the nutritive value of sweet pepper is high especially for vitamin A and vitamin C (Roy et al. 2011).

Organic manures refer to plant and animal waste that undergo decomposition. Farmyard manure (FYM) is a mixture of decomposed dung and urine from farm animals, along with litter and other materials from their fodder. On average, well-decomposed FYM contains 0.5% nitrogen (N), 0.3% phosphorus (P_2O_5), and 0.5% potassium (K_2O). It can provide most of the essential soil fertility elements required by crops, although not always in sufficient quantities or the correct proportions. Manure's plant nutrients become available to crops as it decomposes with the help of soil microorganisms.

Vermicompost is another organic fertilizer derived from the excreta of earthworms. It is rich in NPK (nitrogen, phosphorus, and potassium)

and micronutrients like calcium, magnesium, zinc, and manganese. Vermicompost also contains enzymes such as amylase, lipase, cellulase, and chitinase, which continue to break down organic matter in the soil and release nutrients to plant roots. It offers various benefits, including plant growth promotion, nutrient-rich content, beneficial bacteria, and mycorrhizae.

Similarly, poultry manure is the waste generated by chickens and serves as a valuable organic fertilizer. It supplies both macro and micro nutrients essential for the growth, yield, and quality of horticultural crops. Poultry manure is particularly notable for its higher nitrogen and phosphorus content compared to other organic manures, with average nutrient levels of 1.5% nitrogen, 1.5% phosphorus (P_2O_5), and 1.8% potassium (K_2O). Proper handling and management are essential, as exposed poultry manure can lose up to 50% of its nitrogen within 30 days. Additionally, poultry manure plays a crucial role in supporting crop nutrition and maintaining soil physical and chemical properties.

2. MATERIALS AND METHODS

The present investigation was carried out during the year 2022-23 in Kharif season at Research

Farm, Department of horticulture, MGCGV, Chitrakoot, Satna, Madhya Pradesh.

2.1 Techniques of Analysis

2.1.1 Soil sampling

The soil samples were taken from different places of the experimental field with the help of auger from 12-15 cm depth after clearing the surface vegetation. These samples were properly mixed, air dried in diffused sunlight, finally powdered and again mixed thoroughly. A representative sample of 5 g was taken for each analysis and subjected to mechanical and chemical analysis.

2.1.2 Mechanical analysis of soil

The mechanical analysis of sample soil was done with the help of Bouyoucos Hydrometer method (1952) and result so obtained are presented Table 1.

2.1.3 Chemical analysis of soil

The chemical analysis of soil was conducted to determine the percentage of major elements viz., nitrogen, phosphorus, potash, organic carbon, organic matter, pH and electrical conductivity. Nitrogen was estimated by alkaline permanganate method (Subbaih and Asija, 1956). The phosphorus was estimated by Olsen's Colorimetric method (Olsen et al., 1954) and potash was estimated by Flame Photometric method (Jackson, 1958) respectively. The soil organic matter was estimated by Hydrochloric oxidation method as

suggested by Walkey and Black (1934). The pH of soil was determined by pH meter (Elico pH meter model 2.112).

2.2 Statistical Analysis

The data were analyzed in randomized block design as per procedure of Cochran and Cox (1959). Interpretation of results was made on the basis of "F" test and critical difference at 0.05 probability calculated to compare the treatments. The data were presented by way of tables, graphs and photographs.

2.3 Critical Difference

The significance of differences between the mean values of different treatments for different characters was taken by calculating critical differences (CD) at 5% level of significance as follows $CD = SED \times T$

$$CD = \sqrt{2 \times EM_{ss} / r \times t \text{ value}}$$

T = number of error d.f at 5 % level of significance

$$SED = \sqrt{(2Me/r)}$$

Where,

t = table value of "t" at error degree of freedom

r = number of replications

Table 1. Treatment combination

S. No.	Treatment No.	Treatment Combination
1	T ₁	Control (Farmers Practices)
2	T ₂	RDF (Refuse derived fuel) (250:150:150 kg ha ⁻¹) NPK
3	T ₃	FYM 25 t ha ⁻¹
4	T ₄	Vermicompost 10 t ha ⁻¹
5	T ₅	SPNF (Subhash palekar natural farming) (Bijamrit + Jivamrit ++ Mulching)
6	T ₆	50% FYM (Farm yard manure) + 50% RDF (Refuse derived fuel)
7	T ₇	50% Vermicompost + 50% RDF (Refuse derived fuel)
8	T ₈	SPNF (Subhash palekar natural farming) + Panchgavya
9	T ₉	50% FYM (Farm yard manure) + Panchgavya
10	T ₁₀	50% Vermicompost + Panchgavya
11	T ₁₁	50% RDF (Refuse derived fuel) + Panchgavya
12	T ₁₂	50% FYM (Farm yard manure) + 50% Vermicompost

Table 2. Mechanical composition of soil

Soil Properties	Percentage	Method Employed
Sand	61%	Bouyoucos Hydrometer method (1952)
Silt	26%	
Clay	15%	
Textural	Sandy loam	

3. RESULTS AND DISCUSSION

The results of the present investigation was carried out during the year 2022-23 in Kharif season at the Research Farm, Department of Horticulture, MGCGV, Chitrakoot, Satna, Madhya Pradesh, India.. The results of the investigation regarding the influence organic nutrients and N.P.K. on growth and yield of Bell pepper have been presented in tables and wherever required. The result of the experiment has been presented given table.

3.1 Growth Parameters

3.1.1 Plant height (cm)

The height of the plants was measured from the base to the growing tip of the plants at harvest stage. The mean plant height were worked out and expressed in cm.

3.1.2 Number of branches plant⁻¹

The total number of branches per plant at four growth harvest stage were recorded from five observational plants and mean value recorded.

The observation in term of plant height (cm) at harvest of Bell pepper (*Capsicum Frutescens L.*) was statistically analyzed and has been presented in table. The perusal of the table shows that the application of organic nutrient and N.P.K had significant positive effect on plant height (cm) at harvest of Bell pepper (*Capsicum Frutescens L.*). Among the treatments applied, treatment T₁₂ 50% FYM + 50% Vermicompost exhibited significantly maximum plant height (65.15) at harvest, and closely followed by T₄Vermicompost 10 t ha⁻¹ and T₃FYM 25 t ha⁻¹. Whereas the minimum plant height (48.14) at harvest found in T₁ Control (Farmers Practices). Similar results were reported by Pariari and Khan [2], Shiva et al., (2015) and Adhikari et al. [3] (Table 3).

The observation in term of number of leaves per plant at harvest of Bell pepper (*Capsicum Frutescens L.*) was statistically analyzed and has

been presented in table. The perusal of the table shows that the application of organic nutrient and N.P.K had significant positive effect on number of leaves per plant at harvest of Bell pepper (*Capsicum Frutescens L.*). Among the treatments applied, treatment T₁₂ 50% FYM + 50% Vermicompost exhibited significantly maximum number of leaves per plant (148.66) at harvest, and closely followed by T₄Vermicompost 10 t ha⁻¹ and T₃FYM 25 t ha⁻¹. Whereas the minimum number leaves per plant (125.11) was found in T₁ Control (Farmers Practices). Similar results were reported by Pariari and Khan [2], Shiva et al., (2015) and Adhikari et al. [4] (Table 3).

The observation in term of Number of fruit plant-1 of Bell pepper (*Capsicum Frutescens L.*) was statistically analyzed and has been presented in table. The perusal of the table shows that the application of organic nutrient and N.P.K had significant positive effect on Number of fruit plant-1 of Bell pepper (*Capsicum Frutescens L.*). Among the treatments applied, treatment T₁₂ 50% FYM + 50% Vermicompost exhibited significantly maximum Number of fruit plant-1 (10.45), and closely followed by T₄Vermicompost 10 t ha⁻¹ and T₃FYM 25 t ha⁻¹. Whereas the minimum Number of fruit plant-1 (6.25) found in T₁ Control (Farmers Practices). Similar results were obtained by Gopinath et al. [5], Pariari and Khan [2], Shiva et al., (2015), Adhikari et al. [3] (Table 3) [6-10].

The observation in term of fruit length (cm) of Bell pepper (*Capsicum Frutescens L.*) was statistically analyzed and has been presented in table. The perusal of the table shows that the application of organic nutrient and N.P.K had significant positive effect on fruit length (cm) of Bell pepper (*Capsicum Frutescens L.*). Among the treatments applied, treatment T₁₂ 50% FYM + 50% Vermicompost exhibited significantly maximum fruit length (cm) (9.25), and closely followed by T₄Vermicompost 10 t ha⁻¹ and T₃FYM 25 t ha⁻¹. Whereas the minimum fruit length (cm) (6.11) found in T₁ Control (Farmers Practices) [11-13]. Similar results were obtained by Gopinath et al. [5], Pariari and Khan [2], Shiva et al., (2015), Adhikari et al. [3] (Table 3).

Table 3. Results of organic nutrients and N.P.K. on growth and yield of bell pepper

Treatment No.	Treatment Combination	Plant height (cm) at harvest	No. of leaves per plant at harvest	No. of fruit per plant	Fruit length (cm)	Fruit width (cm)	Weight of fruit (g)	Fruit yield (g) per plant	Fruit yield quintal per hectare
T1	Control (farmers practices)	48.14	125.11	6.25	6.11	5.82	78.25	745.52	275.52
T2	RDF(30:60:30 kg per ha) NPK	51.45	138.41	9.88	7.15	6.88	84.52	854.15	301.22
T3	FYM 25 ton per ha	60.63	141.66	10.33	9.05	7.12	89.88	975.55	303.78
T4	Vermicompost 10 ton per ha	61.45	145.45	10.28	9.17	7.28	90.15	978.58	308.11
T5	SPNF (bijamrit + jivamrit ++ mulching)	58.44	137.02	9.71	8.63	6.45	81.46	893.25	288.45
T6	50% FYM + 50% RDF	56.28	132.52	9.36	8.45	6.52	88.15	878.25	291.27
T7	50% vermicompost + 50% RDF	57.08	131.25	9.52	8.22	6.28	85.63	863.25	293.25
T8	SPNF + panchgavya	51.45	128.15	9.11	8.63	6.10	86.14	841.25	280.11
T9	50% FYM + panchgavya	52.15	132.55	9.63	7.82	6.22	82.63	838.52	286.44
T10	50% vermicompost + panchgavya	58.11	136.25	9.25	8.66	6.41	84.52	841.45	282.81
T11	50% RDF + panchgavya	52.11	131.41	9.05	8.14	6.33	80.25	852.11	283.12
T12	50% FYM + 50% vermicompost	65.15	148.66	10.45	9.25	7.33	91.45	980.15	310.45
	F- Test	S	S	S	S	S	S	S	S
	C.D. at	1.758	0.542	0.031	0.103	0.191	0.976	1.282	2.710
	S.Ed. (+)	0.848	0.262	0.015	0.050	0.092	0.471	0.618	3.683

The observation in term of fruit width (cm) of Bell pepper (*Capsicum Frutescens L.*) was statistically analyzed and has been presented in table. The perusal of the table shows that the application of organic nutrient and N.P.K had significant positive effect on fruit width (cm) of Bell pepper (*Capsicum Frutescens L.*). Among the treatments applied, treatment T₁₂ 50% FYM + 50% Vermicompost exhibited significantly maximum fruit width (cm) (7.33), and closely followed by T₄ Vermicompost 10 t ha⁻¹ and T₃ FYM 25 t ha⁻¹. Whereas the minimum fruit width (cm) (5.82) found in T₁ Control (Farmers Practices). Similar results were obtained by Gopinath et al. [5], Pariari and Khan [2], Shiva et al. (2015), Adhikari et al. [3] (Table 3).

The observation in term of Weight of fruit (g) of Bell pepper (*Capsicum Frutescens L.*) was statistically analyzed and has been presented in table. The perusal of the table shows that the application of organic nutrient and N.P.K had significant positive effect on Weight of fruit (g) of Bell pepper (*Capsicum Frutescens L.*). Among the treatments applied, treatment T₁₂ 50% FYM + 50% Vermicompost exhibited significantly maximum Weight of fruit (g) (78.25), and closely followed by T₄ Vermicompost 10 t ha⁻¹ and T₃ FYM 25 t ha⁻¹. Whereas the minimum Weight of fruit (g) (91.45) found in T₁ Control (Farmers Practices). Similar results were obtained by Gopinath et al. [5], Pariari and Khan [2], Shiva et al. (2015), Adhikari et al. [3] (Table 3) [14-16].

The observation in term of Fruit yield (g) plant⁻¹ of Bell pepper (*Capsicum Frutescens L.*) was statistically analyzed and has been presented in table. The perusal of the table shows that the application of organic nutrient and N.P.K had significant positive effect on Fruit yield (g) plant⁻¹ of Bell pepper (*Capsicum Frutescens L.*). Among the treatments applied, treatment T₁₂ 50% FYM + 50% Vermicompost exhibited significantly maximum Fruit yield (g) plant⁻¹ (745.52), and closely followed by T₄ Vermicompost 10 t ha⁻¹ and T₃ FYM 25 t ha⁻¹. Whereas the minimum Fruit yield (g) plant⁻¹ (745.52) found in T₁ Control (Farmers Practices). Similar results were obtained by Gopinath et al. [5], Pariari and Khan [2], Shiva et al. (2015), Adhikari et al. [3] (Table 3).

The observation in term of fruit yield (quintal/hectare) of Bell pepper (*Capsicum Frutescens L.*) was statistically analyzed and has been presented in table. The perusal of the table shows that the application of organic nutrient and

N.P.K had significant positive effect on fruit yield (q ha⁻¹) of Bell pepper (*Capsicum Frutescens L.*). Among the treatments applied, treatment T₁₂ 50% FYM + 50% Vermicompost exhibited significantly maximum fruit yield (q ha⁻¹) (275.52), and closely followed by T₄ Vermicompost 10 t ha⁻¹ and T₃ FYM 25 t ha⁻¹. Whereas the minimum fruit yield (quintal/hectare) (310.45) found in T₁ Control (Farmers Practices). Similar results were obtained by Gopinath et al., [5], Pariari and Khan [2], Shiva et al., (2015), Adhikari et al. [3] (Table 3).

4. CONCLUSION

Among the various levels of organic nutrient and NPK used in the experiment, treatment T₁₂ 50% FYM + 50% Vermicompost for plant height (65.15) at harvest, number of leaves per plant (148.66) at harvest, Number of fruit plant⁻¹ (10.45), fruit length (cm) (9.25), fruit width (cm) (7.33), Weight of fruit (g) (78.25), Fruit yield (g) plant⁻¹ (745.52), fruit yield (quintal/hectare) (275.52), was the best for the maximum growth and yield of chilli under Satna agro-climatic condition when compared with control and the other treatments.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Tiwari SP, Panigrahi HK, Sharma D. Effect of different fertigation levels on morpho-physiological characters and yield of capsicum under greenhouse condition. International Journal of Agricultural Sciences. 2013;1(9):111-113.
2. A Pariari, S Khan. Integrated nutrient management of chilli (*Capsicum annum L.*) in gangetic alluvial plains. Journal of Crop and Weed. 2013;9(2):128-130.
3. Adhikari P, Khanal A, Subedi R. Effect of different sources of organic manure and chemical fertilizers on growth and yield of sweet pepper. Advances in Plants and Agriculture Research. 2016;3(5):00111.
4. Adhikari P, Khanal A, Subedi R. Effect of different sources of organic manure on growth and yield of sweet pepper. Adv Plants Agric Res. 2016;3(5):00111.
5. Gobinath K. Appireddy, Supradip Saha, Banshi L. Mina, Kundu S, Gupta HS. Effect

- of organic manures and integrated nutrient management on yield potential of bell pepper (*Capsicum Frutescens L.*) varieties and on soil properties. *Journal of Agronomy and Soil Science*. 2011;54:127-137.
6. Ahsanur M, Rahman M, Matiur Rahman, MF Begum, M Firoz Alam. Effect of bio compost, cow dung compost and NPK fertilizers on growth, yield and yield components of chili. *International Journal of Biosciences (IJB)*. ISSN: 2220-6655. 2012;2(1):51-55.
 7. Amla Kanwar, SS Paliyal, Majinder Kaur Bedi. Integrated management of green manure, compost and nitrogen fertilizer in a rice-wheat cropping sequence. *CropRes*. 2006;31(2):334-338.
 8. Bade KK, Vidya Bhati, Singh VB. Effect of organic manures and biofertilizers on growth, Yield and Quality of Chilli (*Capsicum annum*) cv. Pusa Jwala. *Int. J. Curr. Microbiol. App. Sci*. 2017;6(5):2545-2552.
 9. Chandra Harish Raturi, Singh Gurpreet Uppal, Singh SK, Kachwaya DS. Effect of organic and inorganic nutrient sources on growth, yield and quality of bell pepper (*Capsicum Frutescens L.*) grown under polyhouse condition. *Journal of Pharmacognosy and Phytochemistry*. 2019;8(1):1788-1792.
 10. Chandraprabha Koshale, Devendra Kumar Kurrey, Lochan Das Banjare. Effect of organic manure and inorganic fertilizer on growth, yield and physiological parameter of chilli (*Capsicum annum L.*). *International Journal of Chemical Studies*. 2018; 6(4):118-122
 11. Hamendra Sikarwar, Mukesh Kumar Seetpal, Nivedita Singh, Paramita Deb. Effect of NPK and Biofertilizers on Growth and Yield of Chilli (*Capsicum Frutescens L.*). *Biological Forum – An International Journal*. 2023;15(2):73-77.
 12. Holebasappa Kumbhar, A Chandini Raj, Hore JK. Effect of Biofertilizers and Inorganic Fertilizers on Growth and Yield of Chilli (*Capsicum Frutescens L.*). *Int. J. Curr. Microbiol. App. Sci*. 2017;6(7):1564-1568.
 13. Hushal B Muradi, Devi Singh, Deepanshu. Effect of organic manure and inorganic fertilizer on growth and yield of chilli (*Capsicum annum L.*). *The Pharma Innovation Journal*. 2023;12(3):4363-4366.
 14. Kumar VS Tomar, CP Sachan. Effect of organic and inorganic fertilizers combinations on plant growth, fruit yield and yield parameters in chilli (*Capsicum Frutescens L.*). *Int. Arch. App. Sci. Technol*. 2020, September;11(3):182-185.
 15. Maind MM, Yadload SS, Bhalerao RV, Thalkari GN. Effect of irrigation and fertilizer levels on growth and yield of chilli (*Capsicum Frutescens L.*). *Int. J. Curr. Microbiol. App. Sci*. 2018;Special Issue-6:1192-1199
 16. Małgorzata Berova, Georgios Karanatsidis, Krasimira Sapundzhieva, Veselina Nikolova. Effect of organic fertilization on growth and yield of pepper plants (*Capsicum Frutescens L.*), *Folia Horticulturae Ann*. 22/1. 2010;3-7.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/113938>