



Issues and Application of Linearized Models in Geomorphic Properties of the Ikpa Watershed, Nigeria

Umo, Ikpong Sunday^{1,2*#}, M. C. Ike¹ and Ojinma, C. Chux¹

¹*Department of Geography and Environment, School of Social Sciences, Alvan Ikoku Federal College of Education, Owerri, Imo State, Nigeria.*

²*Department of Geography and Regional Planning, University of Benin, Benin, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Author UIS write the manuscript, perform the statistical analyses. Authors MCI and OCC carried out the literature search and support data interpretation. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJEE/2019/10i430123

Editor(s):

(1) Prof. Daniele De Wrachien, Professor of Agricultural Hydraulics at the Department of Agricultural and Environmental Sciences, State University of Milan, Italy.

Reviewers:

(1) Aikaterini Karagianni, Aristotle University of Thessaloniki, Greece.

(2) Uwem Jonah Ituen, University of Uyo, Uyo, Nigeria.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/47234>

Original Research Article

Received 28 November 2018
Accepted 04 February 2019
Published 02 November 2019

ABSTRACT

Though more geomorphological researches are being conducted and previous ideas are being reviewed among geographers and earth scientists, only limited attempts are focused on coupling the relationships among geomorphic attributes on the watersheds in the Humid Tropics. These invariably induced new trends and research directions at diverging scales. This paper is guided by three specific objectives: (1) to employ the Shuttle Radar Topography Mission (SRTM) in deriving the morphologic attributes of the Ikpa Watershed (11) to assess the association and effect of discharge on basin texture, infiltration rate, length of overland flow, and rainfall amount in the Ikpa Watershed, Akwa Ibom State. (111) To describe the relationship between discharge, rainfall amount, and morphologic attributes of the Ikpa Watershed. The watershed was stratified into six sub-basins and data (rainfall and discharge) systematically generated for three months each during the dry and rainy seasons; while the topographic attributes were generated indirectly using SRTM, backup with the four topographic maps. The analysis using multiple linear regression yielded a coefficient of 0.986. Also, the R square value of 0.859 implied that 85.9 percent of the proportion of

*Corresponding author: E-mail: umohikpong@yahoo.com;

#<http://orcid.org/000-0002-8424-6889>

variation in discharge amount of the Ikpa watershed is accounted for by the four independent variables. A comparison of the computed ANOVA result of 8.469 with the Table value of 7.7086 implied that variation in mean discharge is attributed to the significant effect of basin texture, infiltration rate, length overland flow, and rainfall amount in the Ikpa watershed. The results of the curve estimate for linear, logarithmic, and quadratic equations suggested normal predictive ability and each significant at 0.05 confidence level. This study concluded that the dominances of gully erosion at the upstream and peak-flood discharge at the middle and downstream areas of the watershed are directly influenced by the rainfall events and morphology of the landform.

Keywords: Linearized models; morphology; rainfall; discharge; watershed properties.

1. INTRODUCTION

Geomorphology is fundamentally concerned with the Earth's surface, the physical processes that act upon the surface, and the manner in which these processes vary across space and time to create distinctive landforms [1]. Over time sequence, studies on landforms and fluvial processes have been subjected to dynamic paradigms in the conceptual, theoretical, philosophical, and methodological understanding of the surficial environment. Such paradigms often reflect in and influence the various research trends and directions prevalence over a period of time as illustrated in [2,3] and other landform experts.

Climate (rainfall and wind) often played critical role in driving the flux of solutes and mass across eco-geomorphic systems during weathering and denudational processes. In context of geography, [4] observed that the research links between climatologists and geomorphologists are not quite strong to elucidate certain ambiguity in their connection; but more work is still needed on fluvial processes and basin form to elucidate the intriguing relationships which may vary across scale [5,6].

The effect of landscape morphology on the lithology, climate, and life had been evaluated with a view to predicting their relationship but [7] suggested that the challenges of predicting surface evolution could be averted using the conservation mass equation. [8] investigated the influence of river discharge on tidal wave propagation with specific attention to residual water level slope, using a one dimensional analytical model for tidal hydrodynamics. Their study proposed an alternative analytical approach for estimating freshwater discharge on the basis of tidal observations along the estuary.

It is obvious that the success of research in this context has the potential of enhancing the

understanding of the role of the climatic variable (rainfall) and discharge in modifying the morphology of the Earth's surface (basin texture, infiltration rate, length of overland flow) and the feedbacks among the outlined attributes. Researches in the domains are rather eclipsed or not attracting the anticipating outcome. However, one of the best quantitative methods is the use of Multiple Linear Regression Models and is essential in predicting morphologic responses to the discharge scenarios in a medium watershed as attested for in [9]. It is capable of enhancing the efficacy of geomorphic disaster management plans and mitigation efforts.

According to Gregory et al. [10] the British Society for Geomorphologists had emphasized the need to understand the changing forms of communication and present challenges of development in geomorphology over half a century. One of the major agitating issues in this research is: What are these challenges confronting recent researches in geomorphology that stimulates trend in the 21st century?

Generally, the United States National Research Council [11] identified nine grand challenges in Earth's surface processes and landforms. These are:

"What does our planet's past tell us about its future? How do geo-patterns on Earth's surface arise and what do they tell us about processes? How do landscapes influence and record climate and tectonics? How does the biogeochemical reactor of the Earth's surface respond to and shape landscapes from local to global scales? What are the transport laws that govern the evolution of the Earth's surface? How do ecosystems and landscapes coevolve? What controls landscape resilience to change? How does the Earth's surface evolve in the anthropocene? How can Earth surface science contribute toward a sustainable Earth surface?" [11].

Although each of the nine grand challenges in Earth surface processes, and geomorphology, in particular, have been identified by (11) and aspects of it re-envisioned in the annual themes of the conferences by the International Geomorphological Association and symposiums of British Society for Geomorphologists as cited in [1], the most cardinal of the issues and trends in geomorphological research that formed the focus of this article are in two critical and high-priority research areas. The two areas are as follows: (1) Quantitative reconstruction of fluvial landform at small scale because at large scale, details are left for specification. (2) Interactions between landforms morphology, climate, and discharge. Each of the identified areas is enriched in scientific investigation and capable of potentially transforming the field of geomorphology. Hence there is a need to quantitatively couple and model the effect of morphology and rainfall on water discharge in the Humid Tropics of the coastal plain sand/alluvium deposits, Southern Nigeria.

1.1 Aim and Objectives of the Study

The vulnerability of the Coastal Plain Sand (Tertiary) and the Alluvium deposits (quaternary) formations to the erosion, landslide, flood and allied geomorphologic hazard could be attributed to the nature of its geology, climate, and local geomorphology. The aim of this study is to evaluate the linearized interactions among the morphologic, rainfall and discharge characteristics in the Ikpa watershed, Nigeria. To achieve the stated aim, the following are specific objectives:

- (1) To employ the Shuttle Radar Topography Mission (SRTM) in deriving the morphologic attributes of the Ikpa Watershed.
- (2) To assess the association and effect of watershed discharge on basin morphology, and climate in the Ikpa Watershed.
- (3) To describe the linearized relationships between discharge, rainfall amount, basin texture, and infiltration rate in the Ikpa Watershed.

1.2 Research Hypotheses

This study is guided by a null research hypothesis which states that: *"Variations in watershed discharge have no significant association and effect on basin texture, infiltration rate, length overland flow, and rainfall amount in the Ikpa Watershed"*.

1.3 The Study Area

The Ikpa Watershed is located within the coastal plain sands deposits of the Tertiary and the Quaternary times. It is a fifth order basin drained by the Great Cross River Basin of Southeastern Nigeria. The watershed composed of two fourth-orders, ten third-order with numerous second and first-order sub-basins. The area is located within the Humid Tropics with the average annual rainfall of approximately 2648 mm, usually with double maxima during July and September. Absolutely, the Ikpa watershed is located between longitude $7^{\circ}46'34.9''$ and $8^{\circ}3'11.9''$, East of Greenwich Meridian and latitudes $5^{\circ}0'3.801''$ and $5^{\circ}16'49.129''$, North of the Equator [12,13]. The basin area covers parts of Ini, Ikono, Ibiono Ibom, Itu, Uruan and Uyo Local Government Areas of the Akwa Ibom State, Nigeria.

Relief of the basin comprises undulating lowland of the coastal plains which form one of the eco-geomorphologic units in the State. The terrain consists of the dissected coastal plains in the middle and Northern sections (e.g. Nkwot, Obotme, Ikot Andem Itam, Ikot Atia, Itu Atai, and Mbiakpan) with an elevation ranging between 100 – 350 meters above mean-sea level with the prevalent steep slopes traversed by ravines and gullies [12,14,15]. At the middle and lower parts (notably in Mbiabam, Use-Abat, Okpoto, Utuat, Ide Uran) of the watershed, there are valleys with the height of 50 – 80 meters above mean sea level, and broad plains sloping gently toward the Cross River channel with elevation less than 50 meters height (Fig. 1).

2. MATERIALS AND METHODS

The study area was demarcated using Shuttle Radar Topography Mission (SRTM). Base on Strahler classification scheme cited in [12,13], two fourth-orders and four third-order sub-basins were selected using stratified sampling techniques for the determination of the morphologic attributes. The mapped area were rectified and geo-referenced in GIS environment with the help of Arc-GIS 10.2 software assigning Universal Transverse Mercator (UTM), World Geodetic System (WGS) dating from 1984 and revised in 2004) and 32N Zone Projection System with ERDAS Imagine 8.5 as applicable in [15,16]. The sampled sub-basins composed {Idim Duem (1), Iyere Stream (2), Akpan Stream (3), Itam Stream (4), Amoor Stream (5), and Iba Oku Stream (6), as depicted in Fig. 1. The mathematical equations for computing each of

the morphologic parameters are summarized on Table 1.

The velocity was measured by means of surface float using cork, for the three segments while the flow velocity was taken and the mean multiplied by 0.85 to overcome errors emanating from the effects of wind and cross-currents as recently emphasized in [9,19]. The formulas developed in Agor cited in [14,19] was adopted and presented thus:

Discharge (Q) = AV. Where A = Cross-sectional Area; V = Velocity

Cross-sectional Area = Total Stream Segment (depth) {a + b + c +d +e}/ Total Stream Segment

Where; a, b, c, d, and e represent average depths of the differential segments.

XY = Total width of the stream at the point of measurement. Velocity (V) = Flow Distant/Time.




Fig. 1. Location and relief of Ikpa watershed

Table 1. Summary of selected morphologic attributes and the mathematical equation

| Morphologic parameters | Mathematical formula (Equation) | Reference |
|-------------------------|--|-----------|
| Basin texture | $Dt = N\mu/P$; Where, $N\mu$ = Number of streams in a given order and P = Perimeter (Km^{-1}) | [17] |
| Infiltration number | $Ir = Dd \times Fs$; Where, Dd = Drainage density (Km) and Fs = Drainage frequency. | [18] |
| Length of overland flow | $Lf = 1/2D$; Where, D = Drainage density (Km/Km) | [17] |

Data generated from distinct sources (SRTM, meteorological station, and direct field measurements of periodic discharge) were analyzed using a set of multivariate statistics. A multiple linear regression model was used to examine the proportion of variance that is associated with the mean discharge on the linear combination of the four independent (basin texture, infiltration rate, length of overland flow, rainfall amount) variables in the watershed because of its predictive ability. The analysis of variance was used a surrogate to test for significant differences in the mean among the dependent and independent variables in the Ikpa Watershed. The rationale was to determine whether significant variations in the mean discharge is attributed to effect of basin texture, infiltration rate, length of overland flow, rainfall amount in the watershed. A t-statistic was used to test for significant difference between discharge and each independent variable. Finally, the relationship between dependent variable (watershed discharge) and each independent variable (basin texture, infiltration rate, length of overland flow, rainfall amount) were analyzed using (linear regression, logarithmic, quadratic) curve estimate. The rationale was to enhanced graphical depiction and estimate of the relationship between mean discharge and each independent variable in the studied watershed. The IBM Statistical Package for Social Sciences (SPSS) version 21.0 was used for data analyses. A test of significant variations and/or relationship was carried out at 0.05 confidence level.

3. RESULTS AND DISCUSSION

3.1 The Effect of Variations in Independent Variables on the Dependent Variable

The effect of mean watershed discharge (dependent variable) variation on the four independent variables was evaluated using a multiple linear regression model and the result summarized on Table 2. A coefficient of multiple determination offers a very high positive effect of 0.986. Also, the adjusted R square sustained the high value of 0.971 which implied that 97.1 percent of the variation in the mean watershed discharge is attributed to the effect of rainfall, infiltration rate, basin texture, length of overland flow. The standard error associated with model is 8.635.

The ANOVA model is employed to examine the level of significant in the mean watershed

discharge on the basin texture, infiltration rate, length of overland flow, and mean rainfall amount. The result summarized on Table 3 showed that the sum of squares is 2526.515. The residual associated with the model yields 74.577 while the total sum of squares of the ANOVA model is 2601.092. Similarly, the mean square offers 631.629 and the residual of the mean square value offers 4.577. The computed ANOVA value is 8.469. The Table value tested at $(0.05)_{1/4}$ offers 7.7086. Considering the results, the null hypothesis is rejected. It is therefore declared that "*Variations in watershed discharge have significant effect on basin texture, infiltration rate, length overland flow, and rainfall amount in the Ikpa Watershed, Akwa Ibom State*". This is an affirmation that increase in discharge pattern is associated with intra-watershed variables (notably rainfall amount, infiltration capacity of the landform, and morphologic attributes that defined the flow pattern within the watershed). However, the external factors to this coupled variable in the model notably vegetation, sub-surface, anthropogenic interferences tend to contribute low proportion of 2.9% (see Table 2). The results is a strong affirmation of similar studies conducted by [9,14] under the same area but with different parameters.

To establish the effect of each independent variable on the mean watershed discharge, a partial regression model is employed. The results of unstandardized coefficients presented on Table 4 reveals a constant negative value of (-45.775) for mean watershed discharge, 0.724 for basin texture, 12.435 for infiltration rate, 61.699 for length of overland flow, and 0.023 for mean rainfall. Similarly, the standardized coefficients yield 0.968 for basin texture, 0.257 for infiltration rate, 0.396 for length of overland flow, 0.099 for mean rainfall amount. The linearized equation is therefore modelled as following:

$$Y = -45.775 + 0.968x_1 + 0.257x_2 + 0.396x_3 - 0.099x_4 \quad (e = 8.636).$$

Basin texture (x_1) exercised the highest positive effect, followed by length of overland flow (x_3), infiltration rate (x_2), while the lowest effect is rainfall amount (x_4).

Furthermore, the test of significant using student t-test reveals 5.444 for basin texture, 0.671 for infiltration rate, 1.081 for length of overland flow, and 0.470 for mean rainfall. The implication of the results is that only the basin texture has a significant positive effect on the mean watershed discharge at 0.05 confidence level. However, the

effect of rainfall, length of overland flow, and infiltration each was not significant at the 0.05 confidence level. However, the model shows strong distinction from the previous findings reached in the area by [9,15,19] possibly due to differences in context, parameters, and methods used by the researchers.

3.2 Curve Estimate of the Relationship between Independent and Dependent Variables

Discounting the effect of distinct independent variables on the mean watershed discharge established on the preceding section, a curve estimate was employed as a surrogate in depicting the pattern of relationship existing between dependent variable and each independent variable. The result of the linear relationship between mean discharge and basin texture is summarized in Appendix with a high positive linear model of 0.927 and the F value of 51.167. The quadratic equation yields 0.930 and the F value of 19.928 both are significant at 0.05 confidence level. The logarithmic model yields 0.535 and the F value of 4.606 express a

moderate predictive ability but not significant at 0.05 confidence level. The detail of constant value and for the linear parameter estimate is shown in appendix A and the graphical relationship depicted in Fig. 2. The logarithmic equation show positive relationship between discharge and basin texture. The quadratic equation reveals a positive increase at the beginning but turn negative at middle, which is an affirmation of the moderate predictive ability expressed in the model.

In considering the relationship between mean watershed discharge and infiltration rate in the Ikpa Watershed using Logarithmic equations offer very valid negative relationships but quadratic equation turned positive with further increase in discharge. It is an affirmation that increase in the amount of water loss through surface infiltration affect the amount of water available on the surface as runoff (discharge) but as soon as the saturation level is attained the relationship turn to negative as depicted in Fig. 2 and appendix B respectively. Holding factors constant, flood hazard tend to reduce if the landform has high infiltration rate and vice versa.

Table 2. Multiple linear model of the combined effect of mean discharge on four independent variable^b

| Model | R | R square | Adjusted R square | Std. error of the estimate |
|-------|-------------------|----------|-------------------|----------------------------|
| 1 | .986 ^a | .971 | .857 | 8.636 |

a. Predictors: (Constant), Mean_Rainfall, Length_Overland_Flow, Basin_Texture, Infiltration_Rate
 b. Dependent Variable: Mean_Discharge

Table 3. ANOVA model of mean watershed discharge on the four independent variables

| Model | | Sum of squares | df | Mean square | F |
|-------|------------|----------------|----|-------------|-------|
| 1 | Regression | 2526.515 | 4 | 631.629 | 8.469 |
| | Residual | 74.577 | 1 | 74.577 | |
| | Total | 2601.092 | 5 | | |

a. Dependent Variable: Mean Discharge.
 b. Independent Variables: Mean_Rainfall, Length_Overland_Flow, Basin_Texture, Infiltration_Rate

Table 4. Partial regression coefficients^a and zero-order correlation coefficients

| Model | Unstandardized coefficients | | Standardized coefficients | T | Sig. | Zero-order correlations |
|----------------------|-----------------------------|------------|---------------------------|-------|------|-------------------------|
| | B | Std. Error | Beta | | | |
| 1 (Constant) | 45.775 | 51.509 | | -.889 | .537 | |
| Basin_Texture | .724 | .133 | .968 | 5.444 | .016 | .963 |
| Infiltration_Rate | 12.435 | 18.519 | .257 | .671 | .624 | .104 |
| Length_Overland_Flow | 61.699 | 57.087 | .396 | 1.081 | .475 | -.005 |
| Mean_Rainfall | .023 | .049 | .099 | .470 | .720 | .143 |

a. Dependent Variable: Mean_Discharge




Fig. 2. The relationships between mean discharge and the four independent variables

The mean rainfall and mean discharge reveals positive relationship but the relationship in the long run turned negative due to prolonged storm events. This often gives rise to flooding and loss of flow direction as depicted in Fig. 2 and appendix C. The result suggests that rainfall is one of the most outstanding determinants of flow frequency, density, and intensity of the rivers in the Tropics and the Ikpa River in particular. Similar pattern is observed between mean discharge and length of overland flow Fig. 2 and Appendix D. An increase in discharge attracts a progressive rise in the length of overland flow based on the Logarithmic model, but quadratic

model depicts a drastic decline toward the end as discharge gradually declines.

4. SUMMARY AND CONCLUSION

The fluvial landforms and processes operating within the context of watersheds are complex yet vary in space and time. Using the SRTM and spatial tools, the Ikpa watershed was delineated, stratified into sub-units based on Strahler scheme cited [12] and morphologic attributes computed using appropriate formulas presented on Table 1. The assessment of variations and association presented on Tables 2 and 3 showed

that variation in the mean watershed discharge was due to changes in the four distinct independent (basin texture, infiltration rate, length of overland flow, rainfall amount) variables selected for study in the Ikpa watershed and was significant at 0.05 confidence level. However, the assessment of the effect of variation in the mean discharge on each independent variable tends to vary ranging from positive to negative relationship, low to high value, and from one variable to another and was modelled accordingly. A test of significant between discharge and each independent variable led to the conclusion that only basin texture was significant at 0.05 confidence level (Table 4).

Similarly, the quadratic equation showed a positive relationship between mean watershed discharge and basin texture. The relationship between mean discharge and infiltration rate using quadratic equation showed an increasing trend while a logarithmic equation depicted a decreasing trend and later an increasing trend. Besides, the relationship between discharge and rainfall using quadratic model showed positive trend while quadratic model showed a positive relationship at the beginning and later a decreasing pattern. Our findings implied that discharge in the Ikpa watershed is strongly related to and triggered by the rainfall, basin texture, infiltration rate, and overland flow which together defined the pattern of geomorphic hazards in the area.

Considering the converging or diverging effects of variations existing among and between mean discharge and various independent variables at the Ikpa watershed, this study suggested the need to extend similar study to other watershed outside the study area with a focus on both linearized and non-linearized models. An adequate and valid knowledge of distinct geomorphic properties in the Humid Tropics are useful in mitigating the effect of geomorphic hazards in the area as induced by diverse geomorphic, geographic, human, and geologic factors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Legleiter CJ, Marston RA. Introduction to the special issue: The field tradition in

- geomorphology. *Geomorphology*. 2013; 200:1–8.
2. Rhoads BL. Beyond pragmatism: The value of philosophical discourse for physical geography. *Geomorphology*. 1999;760–771.
 3. Rhoads BL. Whither physical geography? *Annals of the Association of American Geographers*. 2004;94(4):748–755.
 4. Dadson S. Geomorphology and earth system science. *Progress in Physical Geography*. 2010; 34:385–398.
 5. Brocklehurst SH. Tectonics and geomorphology. *Progress in Physical Geography*. 2010;34:357–383.
 6. Dlugoss V, Fiener P, Van Oost K, et al. Model based analysis of lateral and vertical soil carbon fluxes induced by soil redistribution processes in a small agricultural catchment. *Earth Surface Processes and Landforms*. 2012;37:193–208.
 7. Dietrich WE. Three grand challenges in geomorphology: Rock, climate, and life. Paper Presented at the (Arthur Holmes Medal Lecture) European Geophysical Union. 2011;13.
 8. Cai H, Savenije HHG, Jiang C. Analytical approach for predicting fresh water discharge in an estuary based on tidal water level observations. *Hydrological Earth System Science Discussion*. 2014;11:7053–7087.
 9. Umo IS, Ike MC, Ugwe IG. Dimensionless geomorphometry and discharge in the Ikpa River Basin, Nigeria. *Asian Journal of Geographical Research*. 2018;1(1):1-13. Available:<http://dx.doi.org/10.9734/AJGR/2018/40098>
 10. Gregory KJ, Lane SN, Lewin J, Ashworth PJ, Downs PW, Kirkby MJ, Viles HA. Communicating geomorphology: Global challenges for the twenty-first century. *Earth Surface Processes/Landform*; 2014. [Retrieved online on 15/6/2015] Available:<http://www.researchgate.net/publication/260912336>
 11. National Research Council. *Landscapes on the edge: New horizons for research on Earth's surface*. Washington D C, National Academies Press. 2010;181.
 12. Udosen CE. Gully erosion in Ikpa River Basin: A threshold phenomenon. *Lagos, Time Communication*; 2008.
 13. Ezemonye MN, Umo IS, Ojinma CC, Ike MC, Enwereuzor AI. Geomorphometric parameters of Ikpa River and its

- implications for the planning of fluvial hazards in the northeast of Akwa Ibom State. *Journal of Geography, Environment and Earth Science International*. 2017; 11(1):1-12.
Available:<http://dx.doi.org/10.9734/JGEESI/2017/34674>
14. Umo IS, Ezemonye MN, Ike MC, Enwereuzor IA. Dimensional basin morphometry and discharge in the coastal plains sands of the Ikpa River, Akwa Ibom State, Nigeria. *Journal of Geography, Environment and Earth Science International*. 2018;13(3):1-13.
Available:<http://dx.doi.org/10.9734/JGEESI/2017/38699>
 15. Umo IS, Ibanga OA, Ike MC. Application of geospatial technologies in assessing gully erosion in the Humid Tropics of Eniong Offot, Uyo, Akwa Ibom State. *Asian Journal of Environment and Ecology*. 2018;6(3):1-13.
Available:<http://dx.doi.org/10.9734/AJEE/2018/4/0224>
 16. Hajam RA, Hamid A, Bhat S. Application of morphometric analysis for geo-hydrological studies using geo-spatial technology –A case study of Vishav Drainage Basin. *Hydrological Research*. 2013;4:157.
Available:<http://dx.doi.org/10.4172/2157-7587.1000157>
 17. Horton R. Erosional development of streams and their drainage basins; hydrophysical approach to quantitative morphology. *Geological Society of America Bulletin*. 1945;56:275-370.
Available:[http://dx.doi.org/10.1130/0016-7606\(1945\)56\[275](http://dx.doi.org/10.1130/0016-7606(1945)56[275)
 18. Zavoianu I. *Morphometry of drainage basins (developments in water science)*. New York, Elsevier Science; 1985.
 19. Umo IS. Drainage basin morphometry and discharge in the Ikpa River Basin, Nigeria. Unpublished M. Sc. Thesis, Department of Geography and Regional Planning, University of Benin, Nigeria. 2014;99.

APPENDICES

Appendix A. Model summary and parameter estimates for mean discharge and basin texture

| Equation | Model summary | | | | | Parameter estimates | | |
|-------------|---------------|--------|-----|-----|------|---------------------|--------|------|
| | R square | F | df1 | df2 | Sig. | Constant | b1 | b2 |
| Linear | .927 | 51.167 | 1 | 4 | .002 | -.233 | 1.287 | |
| Logarithmic | .535 | 4.606 | 1 | 4 | .098 | -12.320 | 16.016 | |
| Quadratic | .930 | 19.928 | 2 | 3 | .019 | 2.147 | 1.012 | .004 |

The dependent variable is Mean_Discharge

Appendix B. Model summary and parameter estimates for discharge and infiltration rate

| Equation | Model summary | | | | | Parameter estimates | | |
|-------------|---------------|------|-----|-----|------|---------------------|-------|------|
| | R square | F | df1 | df2 | Sig. | Constant | b1 | b2 |
| Linear | .011 | .043 | 1 | 4 | .845 | .859 | .002 | |
| Logarithmic | .004 | .016 | 1 | 4 | .907 | .957 | -.021 | |
| Quadratic | .318 | .700 | 2 | 3 | .563 | 1.267 | -.045 | .001 |

Appendix C. Model summary and parameter estimates for mean discharge and mean rainfall

| Equation | Model summary | | | | | Parameter estimates | | |
|-------------|---------------|------|-----|-----|------|---------------------|--------|-------|
| | R square | F | df1 | df2 | Sig. | Constant | b1 | b2 |
| Linear | .020 | .083 | 1 | 4 | .788 | 12.533 | .033 | |
| Logarithmic | .044 | .184 | 1 | 4 | .690 | -51.955 | 13.259 | |
| Quadratic | .378 | .912 | 2 | 3 | .490 | -145.725 | 1.254 | -.002 |

Appendix D. Model summary and parameter estimates for mean discharge and length of overland flow

| Equation | Model summary | | | | | Parameter estimates | | |
|-------------|---------------|------|-----|-----|------|---------------------|---------|----------|
| | R square | F | df1 | df2 | Sig. | Constant | b1 | b2 |
| Linear | .000 | .000 | 1 | 4 | .993 | 21.790 | -.714 | |
| Logarithmic | .002 | .006 | 1 | 4 | .942 | 23.751 | 3.038 | |
| Quadratic | .064 | .103 | 2 | 3 | .905 | -50.309 | 289.855 | -271.918 |

© 2019 Umo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
 The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/47234>